


TETRYONICS

The charged geometry of mass-ENERGY-Matter



Foundational Quantum Mechanics

Abraham

ISBN 978-0-9872884-1-7
[Second Edition © 2012]


questions **Q**

FUNDAMENTUM QUANTUM MECHANICA

A answers

[TETRYONICS]

A fundamental re-interpretation of the geometry of quantised angular momentum is required to complete the physics of 'The Standard model'



Everyone is entitled to their own opinions but NO-ONE is entitled to their own facts


Having removed the impossible, anything that remains, however improbable, must be the truth

Mathematics is the language of Physics, and Geometry is its grammar

Philosophiae Naturalis Principia geometricae

"...the scientist makes use of a whole arsenal of concepts which he imbibed practically with his mother's milk; and seldom if ever is he aware of the eternally problematic character of his concepts. He uses this conceptual material, or, speaking more exactly, these conceptual tools of thought, as something obviously, immutably given; something having an objective value of truth which is hardly even, and in any case not seriously, to be doubted. ...in the interests of science it is necessary over and over again to engage in the critique of these fundamental concepts, in order that we may not unconsciously be ruled by them."

[Albert Einstein]



For ANY physical model to be valid it must demonstrate the known facts

Science is born from observation, and the reasoning of known facts in search of underlying truths

In the following pages the true geometry of quantum mechanics is revealed, leading scientific endeavour into new realms of understanding

mass-ENERGY-Matter


The a-priori revelation of Tetryonic theory is that all square mass-energies possess equilateral momenta geometries



The equilateral Quantised Angular Momentum intrinsic to Planck mass-energy momenta produces charged geometries





The quantum mechanics of velocity, quanta, EM fields and mass-Energy-Matter can be fully revealed through their equilateral geometries




A long hidden topology is revealed

Equilateral triangles are the foundational geometry for all mass-ENERGY-Matter topologies and physical Force interactions




SQUARED energies in quantum mechanics are EQUILATERAL geometries



Square

$$\text{area} = s^2 = [100]$$


Circles



$$\text{area} = \pi * r^2$$

$$= \pi * [5.642]^2$$


$$= 100$$



can be created by a number of planar geometries

For a long time it has been assumed by scientists (and mathematicians) that circular [and squared] geometries are the geometric foundation of all physics, leading to a seriously flawed model of particles and forces in quantum mechanics


Tetryonic theory now reveals that quantised equilateral angular momenta creates the foundational geometry of all the mass-Energy-Matter & forces of physics



Equilateral

$$\text{area} = \left(\frac{1}{2} * b\right) * h$$

Triangles




$$b \quad h$$

$$[.5 * 15.197] * 13.160$$

$$= 100$$

Integers

The integers (from the Latin integer), literally "untouched", hence "whole" in Tetryonics it is the basis for the quantum



Viewed as a subset of the real numbers, they are numbers that can be written without a fractional or decimal component

ODD numbers


An odd number is an integer which is not a multiple of two.

$$2n-1$$

$$2n+1$$

ODD numbers
in each level

Bosons have
ODD number
quanta



An odd number, when divided by two, will result in a fraction

EVEN numbers


An integer that is not an odd number is an even number.

2

Photons have
EVEN number
quanta

2n

EM waves are
comprised of EVEN
numbered quanta



An even number is defined as a whole number that is a multiple of two.
If an even number is divided by two, the result is another whole number.

Square numbers


A square number, sometimes also called a perfect square, is the result of an integer multiplied by itself

$$2n-1$$

Quantum levels have ODD number geometries

$$\Sigma 2n-1$$

SQUARE numbers are the sum of successive ODD numbers



Square numbers

In Tetryonics SQUARE numbers are EQUILATERAL geometries


Square roots

A square root of a number is a number that, when it is multiplied by itself (squared), gives the first number again.

$-i$ and $+i$

Against Mathematical convention,
square roots of negative numbers
are real numbers

In Physics
every complex number
except 0 has 2 square roots.




A whole number with a square root that is also a whole number is called a perfect square

Real Numbers

A real number is a value that represents a quantity along a continuous line.

The real numbers include all the rational numbers,



Irrational Numbers

An irrational number is defined to be any real number that cannot be written as a complete ratio of two integers

$+i$ and $-i$


Well known irrational & imaginary numbers in Math are π and i

π

\sqrt{n} *in Tetryonics the SQR of a negative number is the linear momentum of a negative charge EM field* $\sqrt{-n}$

Irrational numbers often occur in mathematics

$\sqrt{-n}$




Tetryonic Colour Code

0
1
2
3
4
5
6
7
8
9




Brown
Red
Orange
Yellow
Green
Aqua
Blue
Indigo
Violet
Black

A colour code is used to indicate the varying quantum levels of the numerous forms of mass-ENERGY-Matter and serves to illustrate relationships between various Physical properties.



Tetryonics uses a colour code that is based on the spectral colours of dispersed White Light




Free Space

A contiguous volume or area
of any regular geometry
that is free, available, or unoccupied

$$\mathbf{E} = 0$$

in any form



A Spatial region
is defined so as to
measure the physics
of mass-ENERGY-Matter
within its confines

Energy moves through Space
in various forms:
radiant energies, Matter etc

Space can be
Cubic, Spherical or
Tetrahedral as defined by
the spatial co-ordinates
used to define the region

Empty Space is defined as a topology whose volume is devoid of Energy

Space-Time co-ordinates



Charge radiates unilaterally in line with its vector linear momentum

Vector forces

The linear vector of divergent Energy forms various co-ordinate systems based on its vector directions per unit of Time

Euclidean planar space-time




Time in Physics is a measure of how long it takes for light to travel 299,792,458 metres from its source in a vacuum

metre

299,792,458 sec

A metre in Physics is the distance light travels in 1/ 299,792,458 of a second from its source

Cartesian rectilinear space time



299,792,458 m
second

Riemannian curvilinear space-time


Spatial geometries

based on the vector speed of Light
form distinct spatial co-ordinate systems
for the measurement of physics

vector
lines
1 Dimensional
velocity
linear metres



Planar
Circles



radial seconds

Forces



2D planar radiant geometry


mass-energies

Energy has an equilateral geometry and forms
Tetrahedral topologies within any
spatial region



3D standing-wave topology


Cubic
volumes



Cartesian co-ordinates

mass-Energy-Matter
and all forces

Spherical
volumes




radial seconds²

c is the natural velocity of light

Energy has a 2D
equilateral geometry
[Euclidian]

Time - as a measure
of divergent energies
has a radial geometry



Equilateral geometry
is the inverse of
radial geometry

3D Matter has a
tetrahedral mass-energy
topology

The scalar spatial geometry of Energy

Velocity

Velocity is the measurement of the rate and direction of change in the position of an object.

It is a vector physical quantity; both magnitude and direction are required to define it.

The absolute scalar value (magnitude) of velocity is speed, a quantity that is measured in metres per second (m/s or ms⁻¹) when using the SI (metric) system.

v Velocity $\frac{m}{s}$




$$\frac{m}{s}$$

Speed is the scalar value of the Distance traveled per unit of Time

$$\vec{v} = \frac{\Delta x}{\Delta t}$$

Velocity is the vector value of the Distance traveled per unit of Time


$$\vec{\frac{m}{s}}$$





is a 3D SPHERICAL SPACE-TIME co-ordinate system

a acceleration $\frac{m}{s^2}$



Acceleration

In physics, acceleration is the rate of change of velocity [dv] over time [dt]

In one dimension, acceleration is the rate at which something speeds up or slows down.

However, since velocity is a vector, acceleration describes the rate of change of both the magnitude and the direction of velocity.

Acceleration has the dimensions [Length]/[Time Squared]
In SI units, acceleration is measured in meters per second squared (m/s²).

$$a = \frac{\Delta y}{\Delta x} = \frac{\Delta v}{\Delta t}$$

In classical mechanics, for a body with constant mass, the acceleration of the body is proportional to the net force acting on it (Newton's second law)

$$F=ma \longrightarrow a=F/m$$

Force


$$kg \frac{m}{s^2}$$

Additionally, for a mass with constant velocity, (ie in an inertial frame) the energy of motion is expressed as its momentum (acceleration causes changes in Energy-momentum)

$$p = kg \frac{m}{s}$$

Quantised Angular momentum

As it is a physical [equilateral] geometry QAM is conservative in any system where there are no external Forces and serves as the foundational geometric source for all the conservation laws of physics



Conservation of Quantised Angular momenta


In QFT, angular momentum is considered to be the rotational analog of linear momentum, In Tetryonics It is revealed to be the equilateral geometry of mass-energy within any defined spatial co-ordinate system

A major re-definition of quantised angular momenta in physics is revealed



classical rotational angular momenta

≈



Tetryonic quantum mechanics

In quantum mechanics, angular momenta is quantised – that is, it cannot vary continuously, but only in ODD number "quantum steps" between the allowed SQUARE nuclear Energy levels

In physics, angular momentum, moment of momentum, or rotational momentum is a conserved vector quantity that can be used to describe the overall state of a physical system.

When applied to specific mass-energy-Matter systems QAM reveals the true quantum geometry and nature of Energy in our universe

$$h \text{ kg } \frac{m^2}{s}$$



Planck's Constant

$$m \Omega \text{ mass x QAM}$$

Normally viewed as an expression of rotational momentum Quantised Angular Momentum [QAM] is in fact a result: of the equilateral geometric quantization of mass-energy

mass - Energy geometry

Newton developed his
Laws using Force
 $p = mv$
(linear momentum)



Gottfried Leibniz first
described Scalar Energy as
the square of velocity

$E = mv^2$
Leibnitz mass-energy equivalence

mass velocity
squared

$kg \frac{m^2}{s^2}$

Planck developed his Heat
Law using quantised mass
 $h = m\Omega$
(Planck Constant)




Planck's equation for heat energy
describes transverse masses
[Bosons]

$E = nhv$
Planck mass-energy equivalence

quantised mass
per second

Planck quanta per second
 $[kg \frac{m^2}{s} \frac{1}{s}]$



scalar energies can be related to velocity-momenta
through mass geometries

mv^2

Leibniz-Newton mass-Energy equivalence

scalar

$m\Omega v^2$

Tetryonic-Planck mass-Energy equivalence

quantised

hv^2

mass-Energy Forms

Scalar/Linear forms



mass x velocity squared


$$\text{kg} \cdot \left[\frac{\text{m}}{\text{s}} \right]^2$$

Energy momentum

Leibniz (and Newton) showed that the Energy of a system can be viewed as a product of its mass x velocity squared or (equally as linear momentum squared), forever linking Energy to velocity through the scalar property of mass

$$mv^2 = E = p^2$$

Energy is the ability to do work in varying forms such as potential, kinetic, & mechanical energies, work, heat, and chemical or electrical energies.



$$mv^2 = E = hv^2$$

The total energy contained in an object is identified with its EM mass, and Energy (like mass), cannot be created or destroyed

Tetryonic reveals mass to be a scalar measurement of quantised [equilateral] energy per unit of Time

$$m [\Omega] v^2$$

Energy is subject to the law of conservation

Quantised form



Planck quanta per second


$$\left[\text{kg} \cdot \frac{\text{m}^2}{\text{s}} \right] \cdot \text{s}^{-1}$$

mass-Energy quanta

Max Planck revealed that energy was not continuous, it was quantized – only certain energies are allowed. Continuous energy is a scalar property of mass-energy and its quantisation is the result of its equilateral geometry

$$E = hv^2$$

In quantum mechanics energy is defined in terms of the energy operator as a time derivative of the wave function



Energy is mass-velocity squared


$$E = mv^2$$

scalar mass

kg



$$m = \frac{E}{v^2}$$



$$\left[\left[\begin{array}{cc} \text{Planck} & \text{quanta} \\ m & \Omega v \\ \text{mass} & \text{velocity} \end{array} \right]^2 \right]$$


linear momentum squared

Everything in our Universe results from the equilateral geometry of quantised mass-energies

Quantised Energy

$$\left[\left[\begin{array}{cc} \text{Planck quanta} & \text{per second} \\ \text{kg} & \frac{m^2}{s} \\ \text{mass} & \text{velocity squared} \end{array} \right] \frac{1}{s} \right]$$

Scalar Energy



Energy is Planck-quanta squared

$$E = hv^2$$

quantised mass

kg $\frac{m^2}{s}$



$$h = \frac{E}{v^2}$$

Note: There is a direct correspondence between Velocity and Planck quanta numbers (ie as velocity varies, the energy quanta varies as well by the square of the linear change)

Linear Momentum

p Momentum $\text{kg} \frac{\text{m}}{\text{s}}$

In classical mechanics, momentum (pl. momenta; SI unit kg·m/s, or, equivalently, N·s) is the product of the mass and velocity of an object (p).

Like velocity, momentum is a vector quantity, possessing a direction as well as a magnitude.

Momentum is a conserved quantity (law of conservation of linear momentum), meaning that if a closed system is not affected by external forces, its total momentum cannot change.

Momentum should be referred to in its specific forms to distinguish it in its various forms [Quantised Angular, Linear, Rotational and quantum/nuclear momentum]

$$p = \frac{h\nu^2}{\nu} = mv$$


Although originally expressed in Newton's Second Law, the conservation of momentum also holds in special relativity and, with appropriate definitions, a (generalized) momentum conservation law holds in electrodynamics, quantum mechanics, quantum field theory, and general relativity.

In relativistic mechanics, non-relativistic momentum is further multiplied by the Lorentz factor.

$$p^2 = E = mv^2$$

Energy can be expressed as the square of linear momentum

$$\text{kg} \frac{\text{m}^2}{\text{s}^2}$$






Just as Tetryonic geometry distinguishes between angular momenta and linear momentum it also distinguishes between linear momentum and the vector velocities it produces




Linear Momentum

$$p = mv$$

Velocity



$$E = pv$$



$$E = p^2$$

SCALAR square root

Scalars are quantities that are fully described by only their magnitude

$$E = mv^2$$

VECTOR square root


Vectors fully describe both the magnitude and direction.



linear momentum is a scalar component of all equilateral mass-energies that produce vector velocities



Velocity-Quanta equivalence




classical vector force

$$mv^2$$

mass
linear velocity
squared

$$\text{kg} \left[\frac{\text{m}}{\text{s}} \right]^2$$

Classically, the Energy of massive bodies was determined using the Newtonian mass-velocity relationship
but most recently Quantum mechanics was developed utilising the Planck's quantised Energy relationship




quantised energy momenta

$$hv^2$$

mass
angular momenta
per second

$$\text{kg} \left[\frac{\text{m}^2}{\text{s}} \frac{1}{\text{s}} \right]$$




The EM mass-Energy relationship can be revealed either by linear or angular momentum analysis

Energy-momentum relationship


The total number of Planck quanta [mass-angular momenta] in any physical system is directly related to the square of its linear momentum [mass-velocity]

$$h = \frac{E}{v^2}$$




Quantised Energy-momentum

$$m = \frac{E}{v^2}$$



Linear Energy-momentum



mass-energy momenta

The omega geometry of Energy produces the direct relationships between Planck's constant-quanta and mass-Energy-momentum of any spatio-temporal co-ordinate system

$$E = m\Omega v^2$$

Quantised Angular Momenta is an equilateral geometry

$$E = pc$$

Linear momentum is a vector Force

$$m\Omega v^2 = E = mv^2$$

mass is a derived physical property relating Energy momenta to Velocity

CHARGE

Charge is a measure of mass.QAM/second
[the equilateral geometry of Energy]
that gives form to all physics



ElectroMagnetic Charge
is a quantum property
resulting from the equilateral
QAM geometry of mass-Energy



The two ElectroMagnetic charge
geometries possible can be modelled
by the flux of electrical energy
in ideal inductive loops

It is a measure of the arrangement of Planck quanta
geometries/topologies within any specific
space-time co-ordinate system

Clockwise inductive
energy flux

1
[1-0]

Positive charged
mass-energy momenta

$$\frac{h\nu}{c^2}$$

$$\text{kg} \frac{\text{m}^2}{\text{s}} \frac{\text{s}^2}{\text{m}^2}$$

$$\text{kg} \cdot \text{s}$$

1.33518 e-20 s

Counter clockwise
energy flux

1
[0-1]

Negative charged
mass-energy momenta

Measuring charge geometries

Charge comes in two types, called Negative and Positive which create the Law of Interaction (historically, the Law of Attraction)

q

quantised Charge



Q

nett Charge




All charged masses seek equilibrium

Charge geometry is velocity invariant



Clockwise Energy flux



anti-Clockwise Energy flux

Historically defined as a physical property of mass-Matter that causes it to experience a force charge is actually the result of the equilateral geometry of qantised mass angular momenta which provides the physical mechanics of mass-ENERGY-Matter differentiation and interaction

The SI unit of quantity of charge is the Coulomb;


$$C \approx \text{mass seconds} \approx \text{kg.s}$$

which can also be expressed as Amp-seconds

EM Field Geometry



mass-energies are 2D radiant EM field geometries



[299,792,458 m/s]



'c' forms a CONSTANT of proportionality for different spatio-temporal co-ordinate systems used to measure mass-ENERGY-Matter



All Matter are 3D EM standing wave topologies



EM fields create 'interaction-at-a-distance'



Electrostatic Matter has opposing 2D KEM fields

Energy quantisation

All mass-Energy-Matter can only have certain integer Energy-momenta.
[mass-Energy in all its forms is QUANTISED]

q

Net quantised angular momenta
[inductive energy flux]
determines Charge

$$m \left[\frac{\Omega}{c^2} \right]$$

$$\text{kg} \frac{\text{m}^2}{\text{s}} \cdot \frac{\text{s}^2}{\text{m}^2}$$

seconds

Transverse quanta
create Quantum Levels

The equilateral geometry of
quantised angular momentum
creates charged masses

Planck's Constant is in fact
mass x equilateral QAM

$$\text{kg} \cdot \frac{\text{m}^2}{\text{s}}$$

q

Net quantised angular momenta
[inductive energy flux]
determines Charge

$$m \left[\frac{\Omega}{c^2} \right]$$

$$\text{kg} \frac{\text{m}^2}{\text{s}} \cdot \frac{\text{s}^2}{\text{m}^2}$$

seconds

Scalar (nett) quanta
create Square Energies


Planck's Constant
can be described in
a number of differing ways

Scalar Energy x Time

$$\text{kg} \cdot \frac{\text{m}^2}{\text{s}^2} \cdot \text{s}$$


Linear momentum x Distance

$$\text{kg} \cdot \frac{\text{m}}{\text{s}} \cdot \text{m}$$



n.Planck quanta per second is Energy


Tetryonic Mnemonics



Energy momentum



Electrical flow



Many physical relationships
 are can be represented with
DELTA mnemonics




Inertial Force



velocity of propagation


UNITS OF PHYSICS




Tetryonic geometry offers a complete geometric understanding of physical terms such as Charge, mass geometries, Energy densities, Matter topologies and spatial impedance along with their roles in physical mechanics

Scalar mass-energies

have an EQUILATERAL geometry
(60°- 60°- 60°)



$$\sin \frac{\pi}{3} = \sin 60^\circ = \frac{\sqrt{3}}{2}$$



$$\cos 60^\circ = \sin 30^\circ = 1/2$$


$$\sin 60^\circ = \cos 30^\circ = \sqrt{3}/2$$

$$\cos \frac{\pi}{3} = \cos 60^\circ = \frac{1}{2}$$

$$\frac{\sqrt{3}}{2} S$$

The Area of an Equilateral triangle


$\frac{1}{2}$ base x height



$$\tan \frac{\pi}{3} = \tan 60^\circ = \sqrt{3}$$

Pythagorean geometry

Energy geometries within Physics including Special Relativity and Lorentz corrections have historically been incorrectly illustrated as having the geometry of right angled triangles



$$E^2 = p^2 c^2 + m_0^2 c^4$$

Generalizing, we see that the square of the total mass-energies is the sum of the components squared. [shown incorrectly formulated in this above equation]


We can see an origin of distance in spacetime relating to velocity in pc in which Energy is subject to Lorentz corrections [v/c]

$$E = pc.$$

Additionally, EM mass can be directly related to the energy content of a body by the velocity of Energy

$$E = mc^2$$

Physics is geometry, one cannot be separated from the other



$$mv^2 = E = hv^2$$

There are three ways to look at geometry – mathematically, verbally, and visually,

Of the three, Visually will be shown to be superior leading to intuitive understandings of Physics, Chemistry, Electrodynamics and Gravitation along with all their related physical attributes

$$6.629432672 \text{ e-34 J}$$


$$\left[\frac{\text{Planck quanta}}{\text{mass} \cdot \text{velocity}} \right]^2$$

$$7.376238634 \text{ e-51 kg}$$

$$mv = p$$

Tetryonic geometry

The source of all the physical relationships of mass-Energy momenta & their constants is the geometry of equilateral Planck Triangles (and all texts must be corrected)



$$F$$

$$\sum \mathbf{F} = \frac{d\mathbf{P}}{dt} = m \frac{d\mathbf{v}}{dt} = m\mathbf{a}$$

Newton's Second law of Motion is based on changes to linear momentum

$$\mathbf{F} = m\mathbf{a} .$$

Tetryonics and Pi radians

Although not historically considered a physical constant, π appears routinely in equations describing fundamental principles of the Universe, due in no small part to its relationship to the nature of the circle and, correspondingly, spherical coordinate systems.

The quantised equilateral geometry of mass-energy momenta is measured in π radians

Using units such as Planck units can sometimes eliminate π from formulae.

Heisenberg's uncertainty principle, which shows that the uncertainty in the measurement of a particle's position (Δx) and momentum (Δp) can not both be arbitrarily small at the same time:

$$\Delta x \Delta p \geq \frac{h}{4\pi} = \frac{\hbar}{2}$$

Einstein's field equation of general relativity:

$$R_{ik} - \frac{g_{ik}R}{2} + \Lambda g_{ik} = \frac{8\pi G}{c^4} T_{ik}$$

The cosmological constant Λ from Einstein's field equation is related to the intrinsic energy density of the vacuum ρ_{vac} via the gravitational constant G as follows:

$$\Lambda = 8\pi G \rho_{vac}$$

Coulomb's law for the electric force, describing the force between two electric charges (q_1 and q_2) separated by distance r :

$$F = \frac{|q_1 q_2|}{4\pi \epsilon_0 r^2}$$

Magnetic permeability of free space relates the production of a magnetic field in a vacuum by an electric current in units of Newtons (N) and Amperes (A):


$$\mu_0 = 4\pi \cdot 10^{-7} \text{ N/A}^2$$

Kepler's third law constant, relating the orbital period (P) and the semimajor axis (a) to the masses (M and m) of two co-orbiting bodies:


$$\left(\frac{2\pi}{P}\right)^2 a^3 = \omega^2 a^3 = G(M + m)$$

and the Gaussian formula for a Normal Distribution:

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-(x-\mu)^2/(2\sigma^2)}$$






Pi radian mass-energy fascia geometries



All Equilateral triangles have internal angles that add up to 180°


Equilateral Triangles can be tessellated in turn forming larger equilateral geometries

SQUARED numbers in physics are EQUILATERAL geometries



$n\pi$ mass-energy geometries

180° π An equilateral triangle of mass-energy momenta has a geometry of π radians 2π 360°



Tetryonic Cardinal Angles

equilateral mass-energy geometries form tetrahedral mass-Matter topologies

EM
Charge

Equilateral
energies

Q

E

Space


Time

M

m

Matter
topologies

mass
geometries




radiant light

in radial spatial co-ordinate systems defined by the speed of light

Charge
ODD π $\left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$
 Bosons & quantum levels
 ElectroMagnetic mass velocity

Kinetic Energies
EVEN π $\left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$
 Photons & EM waves
 ElectroMagnetic mass velocity




ALL Matter topologies stem from tessellated equilateral mass-energies


Matter
4n π $\left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$
 Fermions
 ElectroMagnetic mass velocity

Matter's 4 π mass-Energies are Lorentz invariant to accelerations

mass-energy geometries



Tetryonic geometry



**equilateral quantised angular momenta
is the foundational geometry of all mass-ENERGY-Matter**

The Golden Triangle

Planck's formulation for Energy is imprecise for use in Tetryonics and does not reflect the velocity-momenta relationship inherent in the equilateral geometry of Energy

Planck quantum levels

Energy is gained or lost in whole number multiples of the quantity $h\nu$

$$E = n[h\nu]$$



$$\text{Charge} \quad \text{EM Field} \quad \text{Planck quanta}$$

$$\text{Bosons \& quantum levels} \quad \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

ElectroMagnetic mass velocity


$$3 = 3 \cdot [1]$$

$$9 = 9 \cdot [1]$$

$$25 = 25 \cdot [1]$$

$$h\nu$$

The generalise formulation of Plank's heat law $E = nh\nu$ is now changed to a specific formulation of $E = [\text{ODD}] h\nu$ for transverse quantum levels [Bosons]



mass-energy momenta are geometrically related to velocity

Tetryonic geometry $[n\pi]$ redefines Planck's quantum formulation for heat energies from a generalised equation for 2D energy momenta into a geometric formulation for all mass-energy momenta in Matter

[all equilateral $[\pi]$ geometries contain square number quanta]

Scalar energy quanta

Energy is gained or lost in equilateral geometries as whole number multiples of the quantity $h\nu^2$

$$E = n[h\nu]$$



$$E = n\pi \left[\left[\frac{m \Omega v^2}{\text{mass velocity}} \right] \right]$$

Planck quanta

$$3 = 3 \cdot [1^2]$$

$$9 = 1 \cdot [3^2]$$

$$25 = 1 \cdot [5^2]$$

$$h\nu^2$$

The general formulation of Plank's heat law is also changed to a specific formulation of $E = h\nu^2$ for scalar EM waveforms [ENERGY]

Scalar field geometries

All scalar fields are comprised of Transverse and Longitudinal mass-energy momenta all of which are formed from equilateral quantum geometries

Charge is a conserved force

Energy is a conserved quantity

Bosons are transverse
ODD planck geometries

Energy is scalar
SQUARE planck quanta

$$E = n\pi [[hv]]$$

Planck quanta



$$E = n\pi [[hv^2]]$$

Planck quanta squared




Planck quanta

Energy is gained or lost in whole number multiples of the quantity hv

Scalar energies

Energy is gained or lost in equilateral quanta containing whole number multiples of hv²




All energy is comprised of EM fields

All squared energy geometries have component quantum levels

Squared energy levels in quantum physics

are in fact equilateral Planck mass-energy geometries



Energy quanta defined

Quantum levels

Charge is gained or lost in odd number multiples of the quantity $h\nu$

$$E = n\pi [[h\nu]]$$

Planck quanta


transverse 
ODD quanta



Quantum

The number of Planck quanta in any physical system

$$\nu = f/2$$



Scalar Energies

Energy is gained or lost in equilateral geometries containing whole number multiples of the quantity $h\nu^2$


$$E = n\pi [[h\nu^2]]$$

Planck quanta squared

 scalar
SQUARE quanta



Frequency
EVEN quanta



 Longitudinal

The number of repeating waveform cycles in a system
 $2\nu = f$

Tetryonic field equation

E

$$E = n\pi \left[\left[\underset{\text{mass}}{m} \overset{\text{Planck quanta}}{\Omega} \underset{\text{velocity}}{v^2} \right] \right]$$

electromagnetic energy is a scalar

m

$$n\pi \left[\left[\underset{\text{mass}}{\epsilon_0 \mu_0} \right] \cdot \left[\underset{\text{mass}}{m} \overset{\text{Planck quanta}}{\Omega} \underset{\text{velocity}}{v^2} \right] \right]$$

mass is an inertial constant that relates Force to acceleration

[v-v]

Q

$$\text{Charge } \text{ODD}\pi \left[\left[\underset{\text{ElectroMagnetic}}{\epsilon_0 \mu_0} \right] \cdot \left[\underset{\text{mass}}{m} \overset{\text{Planck quanta}}{\Omega} \underset{\text{velocity}}{v^2} \right] \right]$$

equilateral charge geometry is Lorentz invariant

KE

$$\text{Kinetic Energies } \text{EVEN}\pi \left[\left[\underset{\text{ElectroMagnetic}}{\epsilon_0 \mu_0} \right] \cdot \left[\underset{\text{mass}}{m} \overset{\text{Planck quanta}}{\Omega} \underset{\text{velocity}}{v^2} \right] \right]$$

KEM fields mass-energy geometries are Lorentz variant to accelerations

M

$$4n\pi \left[\left[\underset{\text{ElectroMagnetic}}{\epsilon_0 \mu_0} \right] \cdot \left[\underset{\text{mass}}{m} \overset{\text{Planck quanta}}{\Omega} \underset{\text{velocity}}{v^2} \right] \right]$$

Matter's 4π mass-energy topologies are Lorentz invariant to accelerations

E

$$\frac{E}{c^2}$$

$$\frac{\Omega}{c^2}$$

$$\frac{1}{2} m \left[\frac{v^2}{c^2} \right]$$

$$4n\pi \left[\frac{E}{c^4} \right]$$

Energy momenta geometry



Energy is the total Planck quanta per second



EM mass is scalar energy per unit of Time



Charge is the nett Angular momentum per spatial co-ordinate system



Kinetic Energy is the Electric field mass-energy of Matter moving at v



mass-energy geometry




Matter topology

mass-Energy equivalence




$$m = \frac{E}{v^2}$$

linear momentum
mass velocity





$$mv^2 = hv^2$$

The relationship between scalar mass-energy,
linear momentum and quantised angular momenta
is the result of equilateral geometry



$$h = \frac{E}{v^2}$$

quantised
angular momenta



mass-Energy geometries


Leibniz - Newton

Planck - Einstein

$$p = mv$$

$$mv^2 = E$$

EM mass





Energy

$$m = \frac{E}{v^2}$$

h
Planck's Constant
 $\text{kg} \frac{\text{m}^2}{\text{s}}$

$$\frac{E}{v^2} = h$$

velocity






quanta

$$mv^2 =$$

$$\text{kg} \frac{\text{m}^2}{\text{s}^2}$$

$$\text{kg} \frac{\text{m}^2}{\text{s}} \frac{1}{\text{s}}$$


$$= hv^2$$



velocity squared

quanta squared

There exists an intrinsic geometric relationship between mass-velocity and the quantisation of Energy momenta



Electro-Magnetic mass


EM mass-energy

$$E = mc^2$$

EM mass is related to Matter
through the square of the velocity of light
of $8.987551787 \times 10^{16} \text{ m}^2/\text{s}^2 \text{ [c}^2\text{]}$

EM mass-Matter

$$E = Mc^4$$




Matter's energy quanta

scalar EM mass



$$M = \frac{E}{c^4}$$



3D Matter has 2D mass-energies moving at c
in a 4π standing wave topology
creating a closed volume


quantised mass



$$h = \frac{E}{v^2}$$

Kinetic Electro-Magnetic fields


[The energies of Motion]




$$p^2 = E = Mv^2$$

The Electric field energy in any EM field is equal and orthogonal to the Magnetic field energy

$$p^2 = KEM = Mv^2$$



All Matter in motion possess momenta and kinetic mass-energies in extrinsic KEM fields



These 2D planar fields are subject to Lorentz factor corrections that apply to the KEM field mass-energies of Matter in motion

Charge & Kinetic EM fields

Charges are the result of quantised angular momenta


Kinetic Energies result from Matter in motion

$$mv^2 = E = hv^2$$

$$p^2 = E = m\Omega v^2$$

$$KE = \frac{1}{2}Mv^2$$

$$KEM = Mv^2$$




EM field energies externalised on the fascia of 3D Matter geometries form the physical property known as elementary Charge

[k]EM energies in any region of free space are viewed as either a Kinetic energy field with an associated Magnetic moment or a neutral charge Photon of ElectroMagnetic mass-energy




$$m_0 = \frac{E}{c^2}$$

Rest mass is equivalent to the total quantity of Energy in a body or system (divided by c^2)




2D fields propagate at c
3D Matter moves at v



inertial mass

$$m = \frac{F}{a}$$



Inscribed equilateral triangles are the inverse of any radial-circular spatial geometry circumscribing them

mass must be distinguished from Matter in physics, because Matter is a poorly-defined concept in modern science, and although all types of agreed-upon Matter exhibit properties of mass, it is also the case that there many types of Energy that posses NO Matter topology, such as potential energy, kinetic energies and electromagnetic radiation (photons)

$$n\pi_{\text{mass}} \left[\begin{array}{c} \text{EM Field} \\ \epsilon_0 \mu_0 \\ \text{ElectroMagnetic mass} \end{array} \cdot \left[\begin{array}{c} \text{Planck quanta} \\ m \Omega v^2 \\ \text{mass velocity} \end{array} \right] \right]$$

The term 'massless' must be re-termed 'Matterless' to reflect true physical attributes of mass-energy-momenta

EM mass is a planar measurement of 2D energy per unit of time

Thus, all 3D Matter topologies have charged fascia comprised of 2D scalar of mass-energies, but closed volume 3D topology is not a property of 2D EM mass-energy geometries.

mass

ρ

EM mass is a measure of planar energy density per second

$$\frac{mv^2}{c^2} \quad \boxed{\frac{m \Omega v^2}{c^2}} \quad \frac{hv^2}{c^2}$$

per unit of time

EM mass geometries are subject to Lorentz corrections

velocity QAM quanta

$$\frac{mv^2}{c^2} = \frac{m \Omega v^2}{c^2}$$

per unit of time

Gravitational Matter



$$g = G \frac{M}{r^2}$$

Electromagnetic mass-energy

In physics, EM mass-energy equivalence is the concept that the EM mass of a body is a measure of its energy content

Using this concept, EM mass is a property of all Energy, and Energy is a property of all EM mass, and the two properties are connected by a constant.

Using Tetryonic geometry it can be shown that the constant is the equilateral geometry of QAM thus unifying Classical mechanics and Relativistic mechanics






2D mass-ENERGY geometry is NOT 3D Matter topology

$$m = \frac{E}{v^2}$$

Classical mass

$$m = \frac{E}{c^2}$$

Relativistic mass



radiant 2D mass-energies are planar equilateral energy geometries

standing-wave 3d mass-Matter are tetrahedral energy momenta topologies

Relativity shows that rest mass and rest energy are essentially equivalent, via the well-known relationship $(E=mc^2)$

EM mass-Energy-Matter



$$\left[\frac{\Omega}{c^2} \right]$$

Charge is a measure of the quantised angular momentum of any physical system

Tetryonic Matter planar spatial Impedance Planck quanta



$$T\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

3D topology ElectroMagnetic 2D mass velocity

Planck quanta

$$\left[m \Omega v \right]^2$$

mass velocity



$$1\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

ODD quanta

$$4n\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

TETRYONS

$$2\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

EVEN quanta

EM mass-Energy momenta

Energy

$$\left[\frac{m \Omega v^2}{s} \right]$$

E

6.629432672 e-34 J

mass-Energy


7.376238634 e-51 kg

m

$$\left[\frac{E}{c^2} \right]$$

mass

Planck's Constant
is the 'quantum of Action'



$6.629432672 \times 10^{-34} \text{ J.s}$

mass momenta

quantised angular
momenta

$$\left[\frac{m^2}{s} \right]$$



0.0012 m²/s


Charge-momenta

1.33518 e-20 s

S

$$\left[\frac{\Omega}{c^2} \right]$$

charge



ρ

2D

EM mass-energy geometry
per square metre



ρ

Energy density
per spatial region

$$\tau\pi \left[\begin{matrix} \text{EM Field} \\ \epsilon_0 \mu_0 \end{matrix} \cdot \begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right]$$


3D Matter ElectroMagnetic 2D mass velocity



ρ

3D


EM mass-Matter topology
per cubic metre



mass

m






kg/s



Matter

M

KG/s²



Zero Point Fields [ZPFs]


IDEAL QUANTUM INDUCTORS
(equilateral triangle Energy geometry)

The EM FIELD

Electric flux fields can propagate in any direction
Magnetic fields are always at 90 degrees to Electric fields

Magnetic dipole fields propagate in 2 directions at 180 degrees to each other (bi-directionally) forming North and South poles

Electric flux field energy is directly proportional to the resultant Magnetic dipole field energy and vice versa



The linear Electric field strength is directly proportional to its associated transverse Magnetic field which propagates bi-directionally from & into the bloch wall of the Zero point field



ZPFs are quantum inductive tank circuits (Short-circuited 'IDEAL' inductors with energy)
ZPFs charge energies do NOT oscillate
[The magnetic dipole vector determines charge]



h
Positive Charge ZPF
Nett positive Planck quanta with North-South m-dipole vector



Zero Point Fields consist of Electric and Magnetic (EM) fields propagating at 90 degrees to each other

Magnetic MONOPLES do NOT exist
Energy quanta always form charged Electric fields and dipole Magnetic fields

As localised energy quanta increases (number of ZPFs per time unit) the charge geometry remains the same



h
Negative Charge ZPF
Nett negative Planck quanta with South-North m-dipole vector





$$mv^2 = E$$

Quantised Angular Momentum

is the equilateral geometry of scalar energy momenta per unit of time

$$E = hv^2$$



p
linear momentum
 $kg \frac{m^2}{s}$

Normally perceived as vector rotation about a point
quantised angular momentum is actually
an equilateral gaussian flux geometry



(there is no physical vector rotation component)

h
Planck's Constant
 $kg \frac{m^2}{s}$

Charged mass & Matter in motion

produce differing but related measurements of EM force




Charged masses per second $\frac{C}{S} \approx \text{kg} \approx A$ Charged Matter in motion
Amperes

By definition in Tetryonics the quantised angular momentum of mass in any defined spatial co-ordinate system provides the quantum basis for


$$\text{kg} \frac{\text{m}^2}{\text{s}}$$

charged mass-energies




Vector linear momentum creates CURRENT

Quantised angular momenta creates CHARGE




charged mass $C \approx \text{kg} \cdot \text{s} \approx \text{As}$
Coulombs Amp.sec

Charge, through its energy interactions, creates the geometric scaffolding for ALL 3D EM mass-Energy-Matter geometries



Charged mass geometry



The symmetry of Charge geometries provides a geometric foundation for all mass-Energy-Matter particles and physical forces



'a un-balanced (broken) energy symmetry resulting from the nett geometry of equilateral EM mass-energy quanta'



electrically modelled as anti-clockwise inductive energy flux

Negative Charge Zero Point Field



Positive Charge Zero Point Field

electrically modelled as clockwise inductive energy flux



Elementary and quantum Charges

"All known Fermions have charge topologies that are integer 1/3 multiples of the elementary charge"

It can be shown that the "quantum of charge" is the intrinsic angular momentum of a Planck mass

It then follows that Quasi-particles [Tetryons and Quarks] have charge topologies with nett charges of 0, 4 and 8

Further, it can be shown that the "elementary charge" applied to Leptons and Baryons (12 times quantum charge) is reflective of their nett charged Matter topologies

topology $T\pi$ charge $[+q]$ geometry







All Charged particles and their respective KEM fields can be modelled with ZPF field geometries reflective of their nett Charge topologies

Opposites attract
Similar repel

quantised Charges
(EM energy flux rotation within a ZPF geometry)

Tetryons

4 **0** **4**
[4-0] [2-2] [0-4]



The elementary charge, usually denoted as *e*, is the electric charge carried by a single proton, or equivalently, the absolute value of the electric charge carried by a single electron



Elementary Charges are the nett charge created by mass-energy-Matter topologies




All Charges seek Equilibrium

Sir Issac Newton



(25 December 1642 – 20 March 1727)



Any changes to velocity result in changes to Energy-momenta within a charged geometry


$$\sum \mathbf{F} = 0 \Rightarrow \frac{d\mathbf{v}}{dt} = 0$$

Newton's first law of motion says: "A body maintains the current state of motion unless acted upon by an external force."


$$\mathbf{F} = m\mathbf{a}$$

Inertia is the resistance of any physical object to a change in its state of motion or rest, or the tendency of an object to resist any change in its motion.

It is proportional to an object's mass.





Ideal Inductive loops (ZPF quanta) will oppose ANY changes to their energy levels and consequently, Inertia can be viewed as an outcome of quantum ZPF self-Inductance



Inertia

The moment of inertia plays much the same role in rotational dynamics as mass does in linear dynamics, describing the relationship between angular momentum and angular velocity, torque and angular acceleration, and several other quantities

The difference between the impeding (inertial) linear momenta and the co-linear (supportive) momenta is a result of the nett Planck mass-energy momenta within the charged geometries of Matter & its KEM fields



Planck Quanta

are often defined as the "quantum of action"

h


Planck's Constant

$$\text{kg} \frac{\text{m}^2}{\text{s}}$$

The Planck constant (denoted h), is a geometric constant resulting from QAM in quantum mechanics



Charge is a result of QAM geometry




Planck quanta are the minimum energy geometry possible (equilateral mass-energy momenta)

quantised angular momentum is the result of equilateral geometry



≈



hv

$$m\Omega v^2$$

Planck quanta per second

$$\text{kg} \cdot \left[\frac{\text{m}^2}{\text{s}} \right] \cdot \text{s}^{-1}$$


Planck's constant relates the energy in one quantum (photon) of electromagnetic radiation to the frequency of that radiation

Photons have neutral quanta geometry



E = n.hv


Planck Number



hv
transverse
Bosons



hf
longitudinal
Photons



$$1e19v = \bar{n} = 5e18f$$

quantum number

$$hv \neq hf$$



$$2\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Photon EM Field Planck quanta
ElectroMagnetic mass velocity

to photon frequency
of quantum misconceptions

BOSONS
are the quanta of Charge

$$1\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Boson EM Field Planck quanta
ElectroMagnetic mass velocity

Equating number of quanta
is the source of a number

PHOTONS
are the neutral quanta of EM waves



The Planck constant was first described as the proportionality constant between the energy (E) of a photon and the frequency (v) of its associated electromagnetic wave. [it in fact describes the energy of bosons]

The zero-point energy for a simple harmonic oscillator of frequency f is 1/2 hf

The relation between the energy and frequency is called the Planck-Einstein relationship.



Quantum Inductors

An "ideal inductor" has inductance, but no resistance or capacitance, and will not dissipate energy (until it interacts with other ZPFs or Matter) and forms the basis for all Charge-Parity-Time [CPT] interactions

A ZPF is fixed in either a + or - state [Quantum Inductor circuit]


Its energy flux direction as modelled electrically is relative to the observer's view or the direction of measurement

Charge polarity is opposed on opposite faces of the same quoin [quantum coin]

The direction of inherent energy flux from the perspective of the observer determines ZPF charge polarity

+ ZPF


A positive ZPF can be viewed as a quantum quoin or an quantum 'ideal' inductor with an internalised energy flux that is the opposite of a negative ZPF



POSITIVE
Quantum L circuit
[quantum inductive circuit]

- ZPF

A negative ZPF can be viewed as a quantum quoin or an quantum 'ideal' inductor with an internalised energy flux that is the opposite of a positive ZPF



NEGATIVE
Quantum L circuit
[quantum inductive circuit]

The Quantum Inductor (L) circuit stores energy as EM mass in π geometries, it does not oscillate

Energy received is stored indefinitely until its release via weak interaction [Inductive Magnetic coupling]

The quantum Inductive circuit is a SINGLE charge tri-field inductive energy loop


It does not oscillate energy between two opposing charges its differing energy fields are the result of its equilateral QAM geometry storing Electric energy in its E field, and Magnetic energy in its M field

The direction of the QAM flux that models inertia is relative to the observer

Quantum Inductors and EM energy Levels



coupled ODD# mass-energies form SQUARE energy geometries



quantum Inductors

Scalar Energies

[1]
Single Quantum Inductor

[1+2]
Coupled
Quantum Oscillator

[1+2]
A Coupled Inductance
in Parallel with
Two Series Inductors
[3+4]


E

[1+2] [3+5] [4+6]
Three coupled Inductances
in Parallel with
Three Series Inductors
[7+8+9]

All quantum levels form inductive magnetic dipole bases
[Weak Force]

Quantum energy levels can be viewed as various combinations of:

Coupled Inductances



Parallel Inductances


$$L_{total} = \frac{1}{\frac{1}{L_1} + \frac{1}{L_2} + \dots + \frac{1}{L_n}}$$


Series Inductances

$$L_{total} = L_1 + L_2 + \dots + L_n$$

The Golden Triangle

Charged mass-energy geometries






Individual equilateral Planck quanta combine to form larger EM mass-energy geometries with the nett Charge being determined by the scalar arrangement of Positive and Negative quanta within the resulting Tetryonic geometry or topology



$$E = n\pi \left[\left[\frac{m\Omega v}{000} \right] \right]$$




Quantum Energy Levels

Within any spatial co-ordinate system the energy density of a region can increase or decrease geometrically creating the familiar squared quantum energy levels of quantum physics



- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8



Magnetic Vectors

Intrinsic Magnetic vectors
are transverse to E fields



External Magnetic fields
are termed B fields

Magnetic vectors can be modelled
geometrically or electrically
through energy field fluxes
[quantised angular momenta]

Opposite charges moving in opposing directions
produce co-directional Magnetic vectors




Within a Magnetic dipole
energy 'flows' South to North
Magnetic Vector A

External to a Magnetic dipole
energy 'flows' North to South




Opposite charges moving in the same direction
produce Magnetic vectors in opposition

Charge Fields



EM energy fluxes in a Positive ZPF flows are electrically modelled as Clockwise (from North to South)



The EM flux directions of Charges can be modelled vectorially with Electric and Magnetic vectors



EM energy fluxes in a Negative ZPF are electrically modelled as counter-Clockwise (from North to South)

All rotational planck energy fluxes can serve as models for the nett quantised angular momenta of any mass-energy geometry

Polarised
Electric and Magnetic fluxes in ElectroMagnetic fields arise from intrinsic quantised angular momentum



EM fields

are the combined Electric & Magnetic fields resulting from the quantised angular momentum of mass-energy in any region of free space



$$\epsilon_0 \mu_0 = \frac{1}{c^2}$$

EM Permittivity-Permeability is a measure of how much resistance is encountered when the quantised angular momenta of EM energies form an electro-magnetic field in a vacuum


Celeritas = 299,792,458 $\frac{m}{s}$

$$c_0 = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$$

EM field Permittivity-Permeability

$$\epsilon_0 \mu_0 = 1.112650056e-17 \frac{s^2}{m^2}$$

$$\left[\frac{A^2 s^4}{kg m^3} \cdot \frac{kg m}{A^2 s^3} \right]$$



m	=	nπ	[ε₀μ₀]	[mΩv²]
mass		charged	Spatial geometry	Energy density

Electric Constant = 8.85418785 e-12 $\frac{F}{m}$ $\frac{A^2 s^4}{kg m^3}$


$\epsilon_0 = \frac{1}{\mu_0 c^2}$

The permittivity of empty space, equal to 1 in centimeter-gram-second electrostatic units and numerically, to 8.854 × 10-12 farad per meter in International System units, where c is the speed of light in meters per second. Symbolized ε₀.

Magnetic Constant = 1.25663706 e-6 $\frac{H}{m}$ $\frac{kg m}{A^2 s^2}$

$k_0 = \frac{1}{4\pi\epsilon_0}$

A measure of the degree to which molecules of some material polarize (align) under the influence of an electric field; symbol k₀, units F/m (farads per metre).




EM field Permittivity

The Electric constant, commonly called the vacuum permittivity, or permittivity of free space, relates the units for electric charge to mechanical quantities such as length and force.


The name Vacuum Permittivity is a misnomer and should be replaced with the correct term EM field Permittivity

The strength of Electric fields is determined by the Electrical Permittivity Constant

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$$



$$\epsilon_0 = \frac{1}{\mu_0 c^2}$$



"Ampere's Law states that for any closed loop path, the sum of the quantities (B.ds) for all path elements into which the complete loop has been divided is equal to the product of μ_0 and the total current enclosed by the loop."

$$k = \frac{1}{4\pi\epsilon_0}$$


The permittivity of empty space, equal to 1 in centimeter-gram-second electrostatic units and to $107/4\pi c^2$ farads per meter or, numerically, to 8.854×10^{-12} farad per meter in International System units, where c is the speed of light in meters per second.

Gauss' Law:


"The total of the electric flux out of a closed surface is equal to the charge enclosed divided by the permittivity"

This applies equally to any geometry chosen to tessellate a surface area

Superpositioned E fields gives rise to Coulomb Forces



Positive Charge Electric Field




Negative Charge Electric Field

Electric permittivity Fields

Negative externalised Planck quanta
(Counter-clockwise energy flux)



Polar view




**Negative
E-fields**

Coupled same charge ZPFs
have **neutralised** Magnetic fields

*In Electro-statics superpositioned E fields
with interactive energy momenta
are the interactive mechanism
for Coulombic forces*

Electrostatic Particles
in motion have Kinetic energies
resulting in Magnetic moments



Positive externalised Planck quanta
(Clockwise energy flux)



Polar view




**Positive
E-fields**


NEGATIVE electrostatic E fields
attract positive charges and
repel negative charges

ϵ_0

POSITIVE electrostatic E fields
attract negative charges and
repel positive charges



**Opposites Attract
Similar Repel**



Vectorial momenta forces in EM fields are bi-directional
due to the energy-momenta quanta comprising them

The currently stated 'standard' premise of Electrical Energy flowing from Positive to Negative is misleading (as Energy also flows from Negative to Positive at the same time)


EM field Permeability

The permeability of free space, also called absolute permeability.

The name Vacuum Permeability is a misnomer and should be replaced with the correct term EM field Permeability

The magnetic constant has the value of $4\pi \times 10^{-7}$ henry per meter.

The strength of Magnetic fields is determined by the Magnetic Permeability Constant




F

The magnetic field is most commonly defined in terms of the Lorentz force it exerts on moving electric charges.

The magnetic field generated by a steady current (a constant flow of electric charges in which charge is neither accumulating nor depleting at any point) is described by the Biot-Savart law

$$\mu_0$$

$$B = \mu_0 H$$



Positive Magnetic Moment

A measure of the degree to which molecules of some material polarize (align) under the influence of an electric field.

Measured in units of units H/m (Henries per metre).

There are NO magnetic monopoles [not under any condition]



Negative Magnetic Moment

Magnetic permeability Fields


Coupled opposite charge ZPFs produce **neutralised** Electric fields



In Magneto statics superpositioned M fields with interactive energy momenta are the interactive mechanism for Lorentz forces




Magneto-static Particles have enhanced Magnetic moments



Ampere's law


Current flowing through a wire will create a magnetic field



magnetic field forces are orthogonal to electric Coulomb forces

Lorentz forces

charged particles in motion are subject to forces due to external EM fields



Vectorial momenta forces in EM fields are bi-directional due to the energy-momenta quanta comprising them

**Opposites Attract
Similar Repel**

NORTH M-fields
attract south M-fields and
repel north M-fields.

μ_0

SOUTH M-fields
attract north M-fields and
repel south M-fields.

Magnetic fields can only exist in conjunction with Electric fields & Electric fields can only exist in conjunction with Magnetic fields

Electro-static fields

Similar charge electric dipole pairings create 'neutralised' Magnetic dipoles

There are no such things as purely Electric or Magnetic fields

Where Energy couples with equal and opposite vectors static KEM fields are created otherwise know as EM fields


All energy fields are Electro-Magnetic in nature a direct product of their equilateral geometry

'Neutralised' electric dipole pairings create Magnetic dipole fields

Magneto-static fields



v=0




Fields of Force



Charged EM field geometries

Opposite charge fields
can produce neutral E-fields
(with magnetic moments)




Electrostatic charged matter generate
charged energy fields around them

Moving charged particles generate
Kinetic energy & Magnetic moments



16
[72-56]

Positive charge electrostatic fields
attract Negative charges
repel Positive charges




*Charged electrostatic fields
accelerate charged particles
vectorally dependent on their quantum
charge mass-energy momenta field geometries*



16
[56-72]

Negative charge electrostatic fields
attract Positive charges
repel Negative charges




E field acceleration of charged particles


Electric fields can accelerate charged particles within their field geometry dependent on the particle's net charged mass-Matter topology

Opposites ATTRACT

Similar REPEL



Positive E fields accelerate Negative charges towards their source



Particle acceleration increases as it approaches the E field source


Positive E fields accelerate Positive charges away from their source



Similar REPEL



Negative E fields accelerate Negative charges away from their source




Negative E fields accelerate Positive charges towards their source



Opposites ATTRACT


The charge quanta within Neutral particles are affected equally by Electric fields

ZPFs can combine to form
4 distinct static EM fields



Electro-static fields

ElectroMagnetic fields



Magneto-static fields


Similar REPEL

[QAM momenta vectors in opposition]


Opposites ATTRACT

[QAM momenta vectors in same direction]

'THE LAW OF INTERACTION'



'FORCES OF INTERACTION'



Positive electrostatic fields

divergent
positive
momenta

convergent
negative
momenta




$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \hat{r}$$

Negative electrostatic fields

convergent
positive
momenta

divergent
negative
momenta



$$\nabla \cdot E = \frac{\rho}{\epsilon_0} = 4\pi k \rho$$

Coulomb's Force Law

$$E = \frac{F}{q} = \frac{kQ_{source}q}{qr^2} = \frac{kQ_{source}}{r^2}$$

Forces of Interaction

"The force of electrical attraction or repulsion between two point charges is directly proportional to the product of magnitude of each charge and inversely proportional to the square of distance between them"

attract Negative charged masses
repel Positive charged masses



attract Positive charged masses
repel Negative charged masses



Electro-Magnetic field Lines

The magnetic field at any given point is specified by both its direction and magnitude


Magnetic dipoles can be produced by the coupling of opposite electric charge fields



or

by accelerating charged bodies of Matter which in turn produce associated Kinetic energies
[Neutral Electric field and equal strength Magnetic moment]

Magnetic South Poles have divergent south vectors and convergent north vectors



Interaction between vectorially opposed (convergent and divergent) vectors within Electric and Magnetic fields produces the familiar lines of force and interactions of Electro-Magnetic fields

Every particle of Matter in motion possesses intrinsic Planck M-field dipoles because of their nett charged quanta which combine to produce a nuclear magneton.



3D Magnetic lines of Force are made up of 2D fields of Planck energy momenta

Magnetic North Poles have divergent north vectors and convergent south vectors


The magnetic field lines of permanent bar magnets are the result of the equilateral charge [QAM] geometry of neutralised quanta that form their electrostatic fields

Magnetic field Forces



attractive magnetic forces

Magnetic moments of same charges moving in the same direction




magnetostatic fields



$$\frac{m^2 \cdot \text{kg}}{s^2 \cdot A^2} = \frac{J}{A^2} = \frac{Wb}{A} = \frac{s^2}{F} = \frac{V \cdot s}{A} = \frac{J/C \cdot s}{C/s} = \frac{J \cdot s^2}{C^2} = \frac{m^2 \cdot \text{kg}}{C^2}$$

In physics superpositioned M fields with interactive energy momenta produce Lorentz forces



$$\mathbf{H} = \frac{\mathbf{B}}{\mu_0}$$

Coupled opposite charge EM fields produce Magnetic dipole fields



(17 December 1797 – 13 May 1878)


$$\mu_B$$

Electrostatic charged particles in motion create magnetic moments



repulsive magnetic forces

Magnetic moments of same charges moving in opposite directions



kinetic EM fields

Magnetic field lines

Magnetostatic Dipole

Magnetic field lines were introduced by Michael Faraday (1791-1867) who named them "lines of force"

Michael Faraday



(22 September 1791 – 25 August 1867)

Magnetic South Poles have divergent south vectors and convergent north vectors

External to the dipole field lines run from North to South

Positive Electric fields have divergent positive E vectors and convergent negative E vectors

Negative Electric fields have divergent negative E vectors and convergent positive E vectors


Tetryonic geometry reveals the true source of all EM field lines of force

Internal to the dipole field lines run from South to North

Permanent Magnets can be viewed as neutral charge Electrostatic EM fields

Magnetic North Poles have divergent north vectors and convergent south vectors

Similar to conventional vs electron current flows Magnetic field vectors can be modelled with two directions of 'magnetic flux'



Parallel Magnetic Dipoles

Michael Faraday




(22 September 1791 – 25 August 1867)

James Clerk Maxwell



(13 June 1831 – 5 November 1879)



Magnetic lines of force are continuous and will always form closed loops.

Magnetic lines of force will never cross one another.

Parallel magnetic lines of force traveling in the same direction repel one another.

Parallel magnetic lines of force traveling in opposite directions tend to unite with each other and form into single lines traveling in a direction determined by the magnetic poles creating the lines of force.

Magnetic lines of force tend to shorten themselves.

Therefore, the magnetic lines of force existing between two unlike poles cause the poles to be pulled together.

Magnetic lines of force pass through all materials, both magnetic and nonmagnetic.

Magnetic lines of force always enter or leave a magnetic material at right angles to the surface

Maxwell had studied and commented on the field of electricity and magnetism as early as 1855/6 when "On Faraday's lines of force" was read to the Cambridge Philosophical Society.

The paper presented a simplified model of Faraday's work, and how the two phenomena were related. He reduced all of the current knowledge into a linked set of differential equations with 20 equations in 20 variables. [Quarterions]

This work was later published as "On physical lines of force" in March 1861.

In his 1864 paper "A dynamical theory of the electromagnetic field", Maxwell wrote, "The agreement of the results seems to show that light and magnetism are affections of the same substance, and that light is an electromagnetic disturbance propagated through the field according to electromagnetic laws

Maxwell showed that the equations predict the existence of waves of oscillating electric and magnetic fields that travel through empty space at a speed of 310,740,000m/s.


His famous equations, in their modern form of four partial differential equations, first appeared in fully developed form in his textbook A Treatise on Electricity and Magnetism in 1873.

The specific features of Faraday's field concept, in its 'favourite' and most complete form, are that force is a substance, that it is the only substance and that all forces are interconvertible through various motions of the lines of force. These features of Faraday's 'favourite notion' were not carried on by Maxwell in his approach to the problem of finding a mathematical representation for the continuous transmission of electric and magnetic forces.


Maxwell considered these electric and magnetic forces to be states of stress and strain in a mechanical aether, a notion further advanced by relativity theory with its 'stress energy' tensor math.

Tetryonics reveals lines of Force to be a direct result of the various superpositioned EM field geometries of equilateral mass-energy momenta

Anti-Parallel Magnetic Dipoles




Magnetic Moments



Single ZPFs are
'ideal quantum inductor elements'


1 Electro-static Energies Charges



*All Matter in motion possesses kinetic energies
which are stored as Planck quanta in their KEM fields*


*The charge geometry of KEM fields are reflective
of the interactive component of the charged topology
of the particle in motion*

Each charge geometry
has distinct Magnetic dipole
alignments




which in turn can only create
2 distinct orientations [spins] of
magnetic moments

2 Magneto-static Energies Kinetic Energies



ZPF sets can form inductively coupled
quantum Harmonic Oscillators



Electrostatic particle modeling

m

EM mass-energy
Tetryonic field geometry

*Tetryonic [4np] standing wave charge fields
form electrostatic Particle topologies
[Charged and Neutral Matter]*

Matter
tetryonic Matter topology

M

Non-neutral
nett
Tetryonic quanta

3




12π geometries
12π topologies

Positive
Charge
Particles

Equal numbers
of opposite
Tetryonic quanta

12π EM field geometries

3



12π EM field geometries



3


12π particle topologies

12π particle topologies

Neutral
Charge
Particles

Non-neutral
nett
Tetryonic quanta

3




12π geometries
12π topologies

Negative
Charge
Particles

*All particles in motion create secondary KEM field geometries
[Kinetic energies and Magnetic moments]*

Electro-static Fields

Kinetic EM Fields




The Kinetic Electric & Magnetic energies are contained in an KEM field extending from a charged particle in motion

Charged particles in motion produce a Magnetic moment



Electric & Kinetic Fields




$$p = Mv$$

$$KE = \frac{1}{2}Mv^2$$

$$m = \frac{E}{v^2}$$

$$E = p^2$$

The combined Kinetic & Magnetic moment energies total MV^2



Kinetic Energy field


KEM field created by a Positive charged body




KEM field created by a Negative charged body



Positive charges repelled
Negative charges attracted



Neutralised Electric fields
Enhanced Magnetic moments



Negative charges repelled
Positive charges attracted

EM & KEM force vectors

All mass-energy quanta
are ideal quantum Inductors

The E&M force vectors
create orthogonal
equilateral EM Fields

Electric fields
propagate orthogonally
to the Magnetic dipole field




Magnetic flux external to the
Magnetic dipole flows
from North to South
&

Internal to a Magnetic dipole
it flows South to North

When ZFFs combine to form
a Magnetostatic dipole
they form orthogonal
magnetic vectors

Positive charge can be modelled
electrically with clockwise
EM fluxes [omegas]


Negative charge can be modelled
electrically with anti-clockwise
EM fluxes [omegas]



Positive charge fields of interaction
result from an excess of divergent
positive energy momenta

Equal
energy-momenta in opposition
create static fields

Negative charge fields of interaction
result from an excess of divergent
negative energy momenta



The opposing E-field
energy momenta
creates the Coulombic
Law of Interaction

The opposing M-field
energy momenta
creates the
Lorentz Force

When E field is at Maximum - B field is at Minimum
when B field is at Maximum - E field is at Minimum

EM Forces and ZPFs

ZPF quanta can combine in differing combinations to produce 3 distinct charged sets

Positive Charges



'Neutralised'
Magnetic dipole moment

Magnetic dipole moment



Neutral Charges

Negative Charges



'Neutralised'
Magnetic dipole moment

All EM mass-Energy-Matter & forces can be modelled using Tetryonic geometries



Positive charge particle KE field and magnetic moment

Zero Point Field EM geometry is the foundation for all the EM forces comprising, and acting between particles of Matter


Lorentz force, Lenz's Law, Right/Left Hand rules, etc can all be easily replaced with this simple geometric model



Negative charge particle KE field and magnetic moment

Kinetic EM fields


All Kinetic EM fields resulting from motion have charge field geometries resulting from the charged Matter geometry of the particle in motion



Positive charge particle topologies produce positive KEM fields



KEM field charge geometries do not contribute to the nett charge



Negative charge particle topologies produce negative KEM fields



Negative KEM field geometries viewed from different angles are positive KEM fields

Positive KEM field geometries viewed from different angles are negative KEM fields




As a direct result of the KEM field being a EM field permeating free space the symmetry of EM fields results in KEM field geometries being viewed as having neutral KE fields with a magnetic moment



Point Particles and KEM fields



e^+
12



e^-
12



Charged rest mass-Matter topology

Quarks T[8π]
Leptons T[12π]

**velocity invariant
rest mass-Matter has a
standing wave topology**

Baryons T[20π]

**Charged Leptons at rest are Electric field standing waves
(with neutral Magnetic poles)
KE from motion generates a Magnetic Moment**

Bosons $n\pi$ [ODD]


**Kinetic EM field geometry
is divergent from a particle's
rest Matter topology**

Photons $n\pi$ [EVEN]



Charge

Q
[v-v]



[12π]
rest Matter
topology

[1π]
Kinetic Energy
geometry



KEM fields of Matter in motion



ZPFs geometries can be used to model the KEM fields of charged particles

2D charged EM field geometries create Matter topologies

tetryons [4π]	tetryons [4π]
quarks [12π]	quarks [8π]
leptons [12π]	leptons [12π]
Baryons [36π]	Baryons [20π]
Elements [84π]	Elements [54π]


The mass-energy content of all charged fascia constituting massive particles have momenta that is proportional to the intrinsic velocity [c] of the standing wave

Relativistic mass

rest Matter



Electro-static particles have neutralised magnetic dipoles



Kinetic motion produces Magnetic moments


Spherical point charges do NOT exist




12 [12-0] e^+ +1 Q -1 e^- [0-12] **12** [0-12]

12n

3D Matter is a standing-wave topology resulting from radiant 2D EM geometries
All leptons and quarks both have 12 charged fascia geometries,
[but differing mass-Matter-particle topologies]




All EM fields resulting from Kinetic Energy (motion) radiate outwards (the intrinsic KEM fields contain both Negative and Positive Energy momenta quanta)



Magnetic field around a current carrying conductor

Magnetic field produced by a Solenoid



μ_0

There is an inverse cube relationship between magnetic field strength and magnetic field force with respect to distance from the magnet.

ϵ_0

Biot-Savart Law

is an equation in electromagnetism that describes the magnetic field B generated by an electric current.

The interaction between Magnetostatic fields for which a Magnetic dipole exists obeys the inverse CUBED law of attraction and repulsion

$1/R^3$

$1/r^2$

The Biot-Savart law is used to compute the magnetic field generated by a steady current, i.e. a continual flow of charges, through a wire, which is constant in time and in which charge is neither building up nor depleting at any point

$1/r^2$

0

$1/r^2$

$1/R^2$


$1/r^2$

The interaction between Electrostatic fields for which an Electric field exists obeys the inverse SQUARE law of attraction and repulsion


$$\mathbf{B} = \int \frac{\mu_0 I d\mathbf{l} \times \mathbf{r}}{4\pi |\mathbf{r}|^3}$$

The radial distance between Magnetic dipoles is less than the distance between Electric dipoles

$$\mathbf{B} = \int \frac{\mu_0 I d\mathbf{l} \times \hat{\mathbf{r}}}{4\pi |\mathbf{r}|^2}$$




Close to one pole of a magnet, B field strength resembles the inverse square of Electric force. This is because it behaves as a "unipolar magnetic field"



Its the equilateral geometry of quantised angular momenta that creates chirality in physics

Mirror imaged Planck quanta are NOT identical to each other




A reflection of Horizontal or Vertical axis results in a changed EM dipole orientation in turn signifying an opposite charge ZPF

Quantum Chirality



The equilateral geometry of any EM field or Matter particle is determined by its nett Coulombic charge



Any nominal rotation about an axis results in a re-orientation of the electromagnetic vectors but does not affect any change to charge etc.

Irrespective of orientation or rotation:

Positive charge fields have clockwise inductive flux geometries

Negative charge fields have counter-clockwise inductive flux geometries


Pi radian - energy momenta geometries



mass-energy momenta


Q

nett charge
[v-v]




π

pi radian energy geometries



quantised angular momentum



$$n\pi \left[\begin{array}{c} \text{Planck quanta} \\ m\Omega v \\ \text{energy} \quad \text{mass} \quad \text{velocity} \end{array} \right]^2$$

π radian geometries are comprised of equilateral mass momenta **Ω**



ZPFs and Bosons

ZPFs are Planck energy quantum elements



Bosons are transverse Quantum levels



Quantised angular momentum is the foundation for all Tetryonic mass-ENERGY-Matter geometries




Bosons are the exchange particles for the EM force




ZPFs are the quanta geometry for Charge, Bosons and Energy

Each Quantum level is an ODD number of quanta (Bosons)

Each 'squared' energy field is the sum of the preceding Bosons



The charged quanta in all mass-ENERGY geometries create a NORMAL Distributions



$$\text{ODD } \pi \left[\left[\begin{matrix} \text{EM Field} \\ \epsilon_0 \mu_0 \end{matrix} \right] \cdot \left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right] \right]$$

Bosons ElectroMagnetic mass velocity



Bosons form the geometry of Quantum Levels



$$W^+ = n\pi_{\text{ODD}} \left[[\epsilon_0, \mu_0] \cdot [m\Omega v^2] \right]$$

Force carrier for Positive charge particles

W⁺ Boson

EM field Planck quanta

$$\text{ODD } \pi_{\text{Bosons}} \left[[\epsilon_0, \mu_0] \cdot [m\Omega v^2] \right]$$

ElectroMagnetic mass velocity


CHARGE carrier Bosons are ODD number quanta

Bosons

form unit charge Quantum levels that facilitate EM induction between mass-Energy-Matter

Neutral Z Bosons and Photons are EVEN quanta Bosons

Neutral charge parallelogram geometry EM force carriers



Z⁰ Boson

Neutral Z Bosons can be formed by combining EVEN numbers of W Bosons

They are EVEN numbered alternate geometry Photons

Photons are alternate (diamond) EM geometries formed from Z Bosons

EMField Planck quanta

$$\text{EVEN } \pi_{\text{Photons}} \left[[\epsilon_0, \mu_0] \cdot [m\Omega v^2] \right]$$

ElectroMagnetic mass velocity



Force carrier for Negative charge particles

$$W^- = n\pi_{\text{ODD}} \left[[\epsilon_0, \mu_0] \cdot [m\Omega v^2] \right]$$

W⁻ Boson



Bosons are charge carriers

[the geometric foundation of quantum levels]




W_+ Positive charge carrier

Negative charge carrier W_-




All Bosons have ODD quanta trapazoidal charged EM mass-energy geometries




Boson Frequencies

All Bosons are 1/2 wavelength EM fields with ODD number quanta
They are the geometric basis for transverse EM field Quantum levels


ODD #
Q
Planck quanta
 $n\pi \left[\frac{m\Omega v^2}{\text{mass velocity}} \right]$



EVEN #
Q
Planck quanta
 $2n\pi \left[\frac{m\Omega v^2}{\text{mass velocity}} \right]$



ODD #
Q
Planck quanta
 $n\pi \left[\frac{m\Omega v^2}{\text{mass velocity}} \right]$



EM Field Planck quanta
ODD $\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[\frac{m\Omega v^2}{\text{mass velocity}} \right] \right]$
Bosons ElectroMagnetic

W Bosons are comprised of
ODD number quanta

Bosons are TRANSVERSE Charge carriers



Neutral Z BOSONS and PHOTONS
have differing EM geometries




Photons are LONGITUDINAL
neutral EM force carriers
resulting from Bosons

EM waves are comprised of
EVEN number quanta [Photons]


EM Field Planck quanta
EVEN $\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[\frac{m\Omega v^2}{\text{mass velocity}} \right] \right]$
EM waves ElectroMagnetic

Boson Waveforms



W The Electro-weak force is the result of Bosons interacting along their edge of Permeability

Boson exchange is the basis of Electro-Magnetic Induction & Charge transfer



The Strong Colour Force is the result of Bosons interacting via their their Electric charge fascia

All ODD Ω geometries [bosons] create a quantum of charge

Positive charge carrier

+1


ODD number quanta

All charges are comprised of odd numbered Bosons
[each Boson is a Quantum level]


ODD number quanta

Negative charge carrier


-1



Integral W Bosons form square Energy geometries
Separated energy geometries create an electromotive force



Bosons




Positive Charges have nett positive quantised angular momenta

Positive Bosons



$$ODD\pi \left[\begin{matrix} \text{EM Field} \\ \epsilon_0 \mu_0 \end{matrix} \cdot \begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right]$$


Bosons ElectroMagnetic mass velocity



Bosons are the geometry of Quantum Levels

Negative Bosons


Negative Charges have nett negative quantised angular momenta



Bosons are TRANSVERSE EM fields [levels]


nett Charge is a SCALAR EM field property

Charge



Positive Charge field


Charge is a measure of the nett quantised angular momenta - energy geometry of any given region of Space-time




Negative Charge field

bosons

photon



The vector linear momentum of all bosons & photons determines their directions of propagation




Electromagnetic induction results from the coupling of ODD quanta [charge boson] mass-energy momenta geometries

Photons are coupled dual charge mass-energy momenta geometries

ODD π


$2n\pi$



mass geometries

$$m \quad n\pi \left[\left[\begin{matrix} \text{spatial field} \\ \epsilon_0 \mu_0 \end{matrix} \right] \cdot \left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right] \right] \quad E$$

mass-energy



the distribution of energy momenta in EM fields creates charged fascia

charged planar mass-energy momenta form
radiant 2D equilateral geometries

energy momenta per second

ρ

mass-energy geometries

m

kg/m²

$$\frac{E}{c^2} = m_{\text{mass}} \quad \text{ENERGY} \quad M_{\text{Matter}} = \frac{E}{c^4}$$

ρ

mass-Matter topologies


M

KC/m³

energy momenta per second squared

charged mass-energy geometries form
standing-wave 3D tetrahedral topologies


Matter displacement topologies have internalised strong force fascia partitions




mass-Matter

$$m \quad 4n\pi \left[\left[\begin{matrix} \text{spatial field} \\ \epsilon_0 \mu_0 \end{matrix} \right] \cdot \left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right] \right] \quad M$$

Matter topologies



4
[4-0]



2D

ρ/sec


EM mass-energy per square metre

EM mass geometries

m

kg/m²

0
[2-2]




Matter

EM mass-energies have 2D planar geometries

$$E = mv^2$$

2D EM mass-Energy forms 3D Matter




EM Matter has a 3D Tetrahedral topology

$$E = Mc^4$$

3D Matter is comprised of 2D EM mass-Energies

0
[2-2]



3D

ρ/sec^2

EM mass-energy per cubic metre

Matter topologies

M

kg/m³


Matter is comprised of $4n\pi$ mass-energy geometries forming a 3D standing wave topology

$$4n\pi \left[\left[\begin{matrix} \text{Tetryons} \\ \text{EM Field} \\ \text{Planck quanta} \end{matrix} \right] \left[\begin{matrix} \text{Matter quanta} \\ \text{ElectroMagnetic} \\ \text{mass} \\ \text{velocity} \end{matrix} \right] \right]$$

Matter is anything that has a closed mass-energy topology and displaces a volume (the 3D massive building blocks of quantum particles and atomic elements)

The EM mass-energy quanta of Matter can be measured transversely and longitudinally allowing them to exhibit a number of wave-like and particle-like properties (de Broglie wavelength & Compton frequency)

4
[0-4]





Tetryons - the quanta of Matter

'massless' is a physics mis-nomer as all energy exhibits mass equivalence

ZPF

mass-Energy quanta

m



$$m = n\pi \left[\left[\frac{1}{c^2} \right] \cdot \left[\frac{\text{Planck quanta}}{\text{mass}} \cdot \frac{\text{Energy}}{\text{velocity}} \right] \right]$$

Energy
momenta

TETRYONS
Platonic tetrahedrons
are the foundational
topologies of all 3D Matter

$$\tau\pi \left[\left[\frac{1}{c^4} \right] \cdot \left[\frac{\text{Planck quanta}}{\text{mass}} \cdot \frac{\text{Energy}}{\text{velocity}} \right] \right] = M$$

Energy
Matter




Tetryon


Matter Quantum

M


2D mass-energy geometries
can be combined to form
3D mass-ENERGY-Matter particles




Standing wave mass-energy geometries and Matter topologies



As each charged fascia's E field tries to propagate outwards it interacts with a M field dipole at each apex




$$M = \tau \pi \left[\frac{\text{spatial}}{\text{impedance}} \cdot \frac{\text{Planck quanta}}{\text{mass velocity momenta}} \right] \left[\frac{1}{c^4} \cdot m \Omega v^2 \right]$$



In turn energised M fields supply energy to the E Fields creating a EM standing wave [the EM topology of Matter]




Matter stores EM mass-energies geometries in its 3D Tetrahedral topology




All mass-energy propagate at the speed of light creating linear energy momentum

Tetryonic charge topologies



The 3D Tetryonic volume of Matter is what distinguishes it from 2D EM mass-energies



TETRYONS

[MATTER]


4π

The four EM mass-energy momenta produced by Tetryonic collapse must not be confused with the classical 4 Energy-momenta (which refers to x,y,z co-ordinates & v).


m

4pi radiant mass-energies

Four 2D mass-energy geometries can combine to form via charge interactions 3D tetrahedral Matter topologies



A 3D Tetryon can be 'collapsed' into a 2D waveform radiant four energy momenta EM waveform that conserves all of its 2D mass-energy-momenta



Tetryon Genesis

1. WEAK interaction
Mutual Inductive Coupling
[Magnetic dipoles interact]

ODD π
charged mass-energies

*Non-Zero
nett mass-Energy-momenta
result in Linear momentum*

Electro-static field



[2-0]

[1-1]



Magneto-static field



*Opposing momenta
result in static EM fields*

[1-1]



Magneto-static field

Electro-static field



[0-2]

2. **EVEN π**
radiant mass-energies
2D EM mass-energies
interact to form
3D Matter topologies

*EM wave momenta
can form standing waves
[Matter geometries]*




[4-0]



[2-2]



[2-2]



[0-4]

3. **STRONG** interaction
Electrostatic Matter attract
via Electric charges and
Magnetic dipoles

4n π
mass-Matter geometries




Positive Tetryon



STRONG force




Neutral Tetryon



STRONG force

Neutral Tetryon



STRONG force



Negative Tetryon



All Tetryonic charges seek equilibrium


Positive Tetryon




charged boson geometry




Negative Tetryon



standing-wave Matter topology




radiant mass-energy geometries





Neutral Tetryons




charged boson geometry



standing-wave Matter topology



radiant mass-energy geometries





mass




$$m = n\pi \left[\left[\frac{\text{spatial}}{1/c^2} \right] \cdot \left[\frac{\text{Planck}}{m} \frac{\text{quanta}}{\Omega} v^2 \right] \right]$$

mass impedance mass velocity momenta

radiant mass-energy geometries



Tetryons



$$E = mc^2$$

Energy per second



energy



$$n\pi \left[\frac{\text{Planck}}{m} \frac{\text{quanta}}{\Omega} v^2 \right]$$

mass velocity

Energy per second²

$$E = Mc^4$$



Matter

$$M = \tau\pi \left[\left[\frac{\text{spatial}}{1/c^4} \right] \cdot \left[\frac{\text{Planck}}{m} \frac{\text{quanta}}{\Omega} v^2 \right] \right]$$

Matter impedance mass velocity momenta

standing wave Matter topologies

mass
geometry

$$n\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$


EM Field
ElectroMagnetic

Planck quanta
mass
velocity

mass-ENERGY-Matter

geometries topologies

ALL EM masss-energies have equilateral geometry



2D mass-energy geometries are a property of 3D Matter topologies




Tetryonic Matter is the building block of all Fermions, elements and compounds

Matter
topology

$$T\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

EM Field
ElectroMagnetic

Planck quanta
mass
velocity



Transverse EM mass-energies form bosons [Quantum levels]

Longitudinal EM mass-Energies form Photons

Bosons combine to form SQUARED scalar EM energies




massless is a scientific misnomer

2D [non-Topoligal] EM mass-ENERGY is 'Matterless'


ALL Matter has a 3D topology

Tetryonic Matter


All quantum Matter has a foundational tetrahedral topology as a result of their equilateral mass-energy geometries [not spherical as has been assumed from the math]



Area sphere = $4\pi r^2$
 Surface curvature = $1/r^2$
 Total curvature = 4π



Spheres & Tetrahedra are both 3D Platonic solids with 4π scalar integral Gaussian topologies and physical displacement volumes [Gauss-Bonnet theorem]




area = $\sqrt{3/4} * a^2$



Area Tetryon
 $A = 4 A_0 = \sqrt{3} a^2$
 Vertex curvature = π
 Total curvature = 4π

Dodecyons

while quarks & leptons are comprised of
 12π mass-energy geometries
 they have differing final Matter topologies



anti-strange quark




8π

strange quark



Quarks have octahedral topologies

anti-down quark



anti-charmed quark

quarks

anti-top quark




down quark

electron neutrino



neutrinos



positron neutrino

$$12\pi \left[\left[\begin{matrix} \text{EM Field} \\ \epsilon_0 \mu_0 \end{matrix} \right] \cdot \left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right] \right]$$

ElectroMagnetic mass velocity



anti-τ neutrino

anti-μ neutrino

tau

12π 0

muon

muon neutrino


tau neutrino



Leptons have dodecahedral topologies

anti-up quark

as mass-energy momenta is added to Matter topologies & KEM fields of motion different generations of particles are created



$$12\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$


dodecyon

EM Field Planck quanta
ElectroMagnetic mass velocity

1. anti-up quark
2. anti-charmed quark
3. anti-top quark



net charge
8
[2-10]



charged mass-Matter topology
8π

Quarks have 12 charged fascia - mass-energy geometries


2 Positive 10 Negative

Quarks have octahedral mass-Matter topologies


[-2/3 elementary charge]

down quark

as mass-energy momenta is added to Matter topologies & KEM fields of motion different generations of particles are created




1. **down quark**
2. **strange quark**
3. **bottom quark**



$$12\pi \left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \end{matrix} \right]$$

dodecyon ElectroMagnetic mass velocity




Quarks have 12 charged fascia - mass-energy geometries

4 Positive 8 Negative

Quarks have octahedral mass-Matter topologies

[-1/3 elementary charge]

anti-down quark




$$12\pi \left[\left[\begin{matrix} \text{EM Field} \\ \epsilon_0 \mu_0 \end{matrix} \right] \cdot \left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right] \right]$$

dodecyon ElectroMagnetic mass velocity


as mass-energy momenta is added to Matter topologies & KEM fields of motion different generations of particles are created

1. **anti-down quark**
2. **anti-strange quark**
3. **anti-bottom quark**



net charge

4
[8-4]



charged mass-Matter topology

8π

Quarks have 12 charged fascia - mass-energy geometries


8 Positive 4 Negative

Quarks have octahedral mass-Matter topologies

[+1/3 elementary charge]

up quark

as mass-energy momenta is added to Matter topologies & KEM fields of motion different generations of particles are created



$$12\pi \left[\left[\begin{matrix} \text{EM Field} \\ \epsilon_0 \mu_0 \end{matrix} \right] \cdot \left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right] \right]$$


dodecyon ElectroMagnetic mass velocity

1. **up quark**
2. **charmed quark**
3. **top quark**



nett charge

8
[10-2]



8π

Quarks have 12 charged fascia - mass-energy geometries

10 Positive 2 Negative

Quarks have octahedral mass-Matter topologies

[+2/3 elementary charge]

Charged mass geometries & Matter topologies are the physical basis for differentiating particles

DODECAHEDRALS

Quarks and Leptons are both 12π [dodecadeltahedral] mass geometries (differentiated by their final nett charged Matter topologies)

OCTAHEDRALS


TETRAHEDRALS

Tetryons are the [4π] tetrahedral quanta of Matter

hv

BOSONS

ZPFs are single quanta Bosons



PHOTONS

Photons are dual quanta Bosons


2D mass
[radiant energies]

All mass-energy-Matter geometries are comprised of Bosons containing quantised angular momenta


3D Matter Fermions

[standing-wave energies]

tetryon



tetryon



quark



anti-quark



quark



anti-quark




Lepton

12
[12-0]



ν_e
 ν_μ
 ν_τ

0
[6-6]



ν_e
 ν_μ
 ν_τ

0
[6-6]

neutrino



Lepton

12
[0-12]

1

2/3

2/3

0

0

1/3

2/3


1

Elementary Charges

divergent negative E-fields

electron

convergent negative E-fields




0
[6-6]

divergent negative E-fields

**electron
neutrino**

convergent positive E-fields



Leptons have 12 charged mass-energy fascia geometries

*Repulsive STRONG force
creates Lepton topologies*

12π

**Tetryonic charged fascias creates
the Matter topologies of all Leptons**


*Repulsive STRONG force
creates Neutrino topologies*

Leptons have dodecadeltahedral mass-Matter topologies

divergent positive E-fields

positron

convergent positive E-fields



0
[6-6]

divergent positive E-fields

**positron
neutrino**

convergent negative E-fields




Charged Leptons

$$12\pi \left[\left[\begin{matrix} \text{EM Field} & \text{Planck} & \text{quanta} \\ \epsilon_0 \mu_0 & m & \Omega v^2 \end{matrix} \right] \right]$$

Dodecyon ElectroMagnetic mass velocity

e+
12



1. **positron**
2. **anti-muon**
3. **anti-tau**


Leptons have 12 charged mass-energy fascia geometries
12 positive charge fascia or 12 Negative charge fascia

repulsive dodecyon geometries

lepton topologies

Charged Leptons have dodecadeltahedral mass-Matter topologies
[1 elementary charge]

e-
12



electron



1. **electron**
2. **muon**
3. **tau**


as mass-energy momenta is added to Matter topologies & KEM fields of motion different generations of particles are created

repulsive charge dodecyon geometry




lepton mass-Matter topology

positron



12
[12-0]

+1



12n



$$12\pi \left[\left[\begin{array}{c} \text{EM Field} \\ \epsilon_0 \mu_0 \end{array} \right] \cdot \left[\begin{array}{c} \text{Planck quanta} \\ m \Omega v^2 \end{array} \right] \right]$$

Dodecyon ElectroMagnetic mass velocity

1. **positron**
2. **anti-muon**
3. **anti-tau**

Neutral Leptons


$$12\pi \left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & [m\Omega v^2] \end{matrix} \right]$$

Dodecyon ElectroMagnetic mass velocity


ν^-
0



nett charge
0
[6 6]



charged
mass-Matter
topology
 12π



1. **electron neutrino**
2. **muon neutrino**
3. **tau neutrino**

Neutrinos have 12 charged mass-energy fascia geometries
6 Positive & 6 Negative

repulsive dodecyon geometries


lepton topologies

Neutrinos have neutral dodecadeltahedral mass-Matter topologies
[0 elementary charge]

ν^+
0



nett charge
0
[6-6]



charged
mass-Matter
topology
 12π



1. **positron neutrino**
2. **anti-muon neutrino**
3. **anti-tau neutrino**

as mass-energy momenta is added to
Matter topologies & KEM fields of motion
different generations of particles are created


electron



repulsive charge dodecyon geometry



lepton mass-Matter topology



$$12\pi \left[\left[\begin{matrix} \text{EM Field} \\ \epsilon_0 \mu_0 \end{matrix} \right] \cdot \left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right] \right]$$



Dodecyon ElectroMagnetic mass velocity

1. electron
2. muon
3. tau



Leptonic Quantum levels




[KEM field mass-energy geometry]



v^2	n
Total quanta	Electron quantum level
1.2e20	1
4.8e20	4
1.08e21	9
1.92e21	16
3.00e21	25
4.32e21	36
5.88e21	49
7.68e21	64




Illustrative schema only: All Leptonic quantum levels have the same equilateral KEM geometry as the compton frequency of the KEM field increases the wavelengths of the quanta decrease



Electron energy levels

As the total Energy content of particles increase the Quantum levels of their KEM fields increase as well, in turn this is reflected in their associated Compton Frequencies and deBroglie wavelengths





All Tetryonic geometries [EM fields & Matter] absorb and release energy exponentially in equilateral Quantum steps

-1
EM electron (spherical model)


The ratio of the area of a sphere
 $A = 4\pi r^2 = \pi d^2$
 to
 the area of a 3 Tetrahedron Lepton shape
 $A = 4A_0 = \sqrt{3}a^2$
 is
2.418399152

12 [0-12]
Tetrahedron electron (tetryonic model)

The spherical model of an electron's topology is wrong




Leptronic Kinetic EM Fields



Kinetic field modelling

RE

Tetryonic
Positive Charge
KEM field geometry



Mv^2


2D
Kinetic
Energy
field

$E = mc^2$

3D
rest
mass-Matter
topology

Positron

The KEM fields of all Matter particles in motion can be modelled using ZPF EM geometries



Similar charges moving in the same direction produce attractive magnetic dipoles

Opposite charges moving in the same direction produce repulsive magnetic dipoles


F_A

Opposite charges moving in different directions produce attractive magnetic dipoles

Similar charges moving in opposing directions produce repulsive magnetic dipoles

Kinetic EM field geometry determines the Magnetic Moments of all particles in motion

Tetryonic
Negative Charge
KEM field geometry



RE

Mv^2

2D
Kinetic
Energy
field

$E = mc^2$

3D
rest
mass-Matter
topology

Electron

The KEM E-field is reflective of the nett particle charge and forms a directional vector of motion

Lepton Families & Generations



Charged Matter topologies in motion creates Kinetic EM field geometries


Charged mass-energy geometries form elementary particle topologies

$$12\pi$$

$$KEM = Mv^2$$

Total relativistic mass-energies [rest Matter + KEM]


All high mass-energy particles 'decay' into low mass-energy particles by emitting photons



Note: All Lepton geometries are the same size - only the energy density of the EM field changes

Lepton Helicity and Chirality

Particle Spin and Handedness is always referenced with respect to the direction of nett total system linear Momentum.




Parallel spin
Creates higher energy magnetic moment


Anti-Parallel spin
Creates lower energy magnetic moment

All Lepton SPINs are relative to the Nuclear Magnetron


Nuclear Magnetron dipole




Particle Chirality



Particle Helicity



Lorentz Force and Transformation



'Point Charges'

The only true 'point charges' are Zero Point Fields



Positive Particles



Negative Particles

Static charge particles have neutral M Fields
Charges in motion have magnetic moments

Relativistic mass-energy Model

Kinetic Energies

velocity related mass-energy geometry


Magnetic moment

12 [12-0]

e^+

1.2e20

Invariant rest Matter topology



Relativistic mass-energy Model

Kinetic Energies

velocity related mass-energy geometry

Magnetic moment

12 [0-12]


e^-

1.2e20

Invariant rest Matter topology

It is the Kinetic EM field geometry of charged Matter topologies in motion that produces Magnetic Moments

[not a relativistic distortion of spherical charge topologies]




Mesons

are 24π subatomic mass-Matter particles composed of one quark and one antiquark


$$24\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Mesons
EM Field
Planck's quanta
ElectroMagnetic
mass
velocity




prevented by strong repulsive forces

24π charged mass-energy fascia geometries



Short lived combinations

14π final mass-Matter topologies








prevented by strong repulsive forces

Tetryonics 09.01 - Mesons

PION decay

24π dual-quark mass-energy geometries

12π lepton mass-energy geometries



24π Mesons $\left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$
ElectroMagnetic mass velocity

24π dual dodecyon geometries
Despite having 24π geometries Mesons have same net Charges as 36π Baryons
36π tri-dodecyon geometries

$12\pi \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right] + 12\pi \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$
Lepton - Neutrino generational pairing produced is dependent on the mass-energy levels of the interacting Pions

24π

+

12π

=

36π

mesons

charged mass-energy geometries

Baryons

Short-lived Particles

All particles seek charge equilibrium

Stable Particles

+16
[20-4]



+12
[18-6]



+8
[16-8]



+4
[14-10]



π+

0
[12-12]



π0

0
[12-12]



π0

-4
[10-14]



π-

-8
[8-16]



-12
[6-18]



-16
[4-20]



Pions



8

[10-2]

4

[8-4]



quarks




4

[4-8]

8

[2-10]



P+



Proton

12
[24-12]

Neutron

N0



0
[18-18]

Nuclear strong force fascia bonds
create Baryonic Matter topologies

N0



0
[18-18]

anti - Neutron

anti-Proton

P-



12
[12-24]

All Matter topologies have internalised mass-energy fascia

mesons

mass-Matter topologies

Baryons

14π

+

8π

=


20π

Baryons

Baryons are 36π mass-energy geometries


$$36\pi \left[\left[\begin{matrix} \text{EM Field} & \text{Planck} & \text{quanta} \\ \varepsilon_0 \mu_0 & m & \Omega v^2 \end{matrix} \right] \right]$$

Baryons ElectroMagnetic mass velocity



that result in 20π mass-Matter topologies

Proton




Tri-quark mass-energy geometry

$$36\pi \left[\begin{array}{c} \text{EM Field} \\ \epsilon_0 \mu_0 \end{array} \cdot \begin{array}{c} \text{Planck quanta} \\ m \Omega v^2 \end{array} \right]$$

Baryons ElectroMagnetic mass velocity

Baryon mass-Matter topology




12
[24-12]




20π
charged mass-Matter topology

Proton



Neutron




Tri-quark mass-energy geometry



$$36\pi \left[\begin{array}{c} \text{EM Field} \\ \epsilon_0 \mu_0 \end{array} \cdot \begin{array}{c} \text{Planck quanta} \\ m \Omega v^2 \end{array} \right]$$

Baryons ElectroMagnetic mass velocity


Baryon mass-Matter topology



0
[18-18]






anti-Neutron




Tri-quark mass-energy geometry

$$36\pi \left[\begin{array}{c} \text{EM Field} \quad \text{Planck quanta} \\ \left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \\ \text{Baryons} \quad \text{ElectroMagnetic mass} \quad \text{velocity} \end{array} \right]$$

Baryon mass-Matter topology



anti-Proton




Tri-quark mass-energy geometry

$$36\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

EM Field Planck quanta
Baryons ElectroMagnetic mass velocity

Baryon mass-Matter topology




12
[12-24]



charged mass-Matter topology


anti-Proton

20π



Baryon geometry


Proton



12

Tetryonic Charge

[24-12]



+1
elementary charge
Proton

[36]


Linear-Tri-quark geometries - prior to Strong force interaction

Protons are
GEOMETRIC MIRROR IMAGES
of Neutrons

[18-18]

Tetryonic Charge


0



N^0

0
elementary charge
Neutron

Neutron




Chared particle families

[charged mass geometries & Matter topologies]



1π



4π

4π

12π mass geometries



Quarks

8π Matter topologies


12π

Leptons

12π Matter topologies




12π



24π

14π




36π

20π

Tetryonic Energy levels

$$E = n\pi \left[\frac{\text{Planck quanta}}{\text{mass}} \frac{\text{Energy}}{\text{velocity}} v^2 \right]$$




charged field quantisation

Every charged fascia of a Field or Particle holds squared energy levels [equilateral EM mass-energy geometries & Matter topologies]

The Tetryonic KEM Colour code

As the energy levels of atomic nuclei increase/decrease the KEM fields of leptons bound to them change accordingly




Electron



All energy levels are Tetryonic square number quantas ie 1, 4, 9, 16, 25 etc (1,2,3,4,5)


rest mass-Matter



The total mass-energy for any particle is its rest mass-Matter + Kinetic Energies


The mass-energies of its charge geometry determines its rest mass-Matter

An electron's nuclear kinetic energy level is determined by the Baryons it binds to (or by incident photons)



Nuclei




Nuclear Quantum Level
electron Kinetic Energy Level



As the Nuclei quantum level increases - so does the electron's energy and momenta

$$KEM = 4\pi \left[\frac{\text{EM field}}{\text{Kinetic energies}} \frac{\text{Planck quanta}}{\text{ElectroMagnetic mass}} \frac{\text{Energy}}{\text{velocity}} v^2 \right]$$

tri-quark charged topology combinations



Quarks seek charge equilibrium by forming Baryonic Matter topologies as a result of charge interactions [STRONG nuclear Force]

- UP } gen 1
- DOWN } gen 1
- CHARMED } gen 2
- STRANGE } gen 2
- TOP } gen 3
- BOTTOM } gen 3

$$\left[\begin{matrix} +8 & +8 & +8 \\ [10-2] & [10-2] & [10-2] \end{matrix} \right] 24 \text{ } \left(\begin{matrix} + \\ \oplus \end{matrix} \right) 2 \text{ } \begin{matrix} 2/3 & 2/3 & 2/3 \end{matrix}$$

$$\left[\begin{matrix} +8 & -4 & +8 \\ [10-2] & [4-8] & [10-2] \end{matrix} \right] 12 \text{ } \left(\begin{matrix} + \\ \oplus \end{matrix} \right) 1 \text{ } \begin{matrix} 2/3 & 1/3 & 2/3 \end{matrix}$$

The particle Zoo is the result of tri-quark mass-energy topologies

$$\left[\begin{matrix} -4 & +8 & -4 \\ [4-8] & [10-2] & [4-8] \end{matrix} \right] 0 \text{ } \left(\begin{matrix} 0 \\ \circ \end{matrix} \right) 0 \text{ } \begin{matrix} 1/3 & 2/3 & 1/3 \end{matrix}$$

$$\left[\begin{matrix} -4 & -4 & -4 \\ [4-8] & [4-8] & [4-8] \end{matrix} \right] 12 \text{ } \left(\begin{matrix} - \\ \ominus \end{matrix} \right) 1 \text{ } \begin{matrix} 1/3 & 1/3 & 1/3 \end{matrix}$$

Twice the number of combinations are possible with the inclusion of anti-matter topologies

$$36\pi \left[\left[\begin{matrix} \text{EM Field} \\ \epsilon_0 \mu_0 \end{matrix} \right] \cdot \left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right] \right]$$

Baryons ElectroMagnetic mass velocity


Any combinations of inter-generational quarks results in differing final mass-energies in the created Baryonic Matter due to energy equalisation

The Particle Zoo

uuu	udu	uud	usu	uts	ubu	suu	sdu	scu	ssu	stu	sbu
uud	udd	ucd	usd	utd	ubd	sud	sdd	scd	ssd	std	sbd
uuc	udc	ucc	usc	utc	ubc	suc	sdc	scs	ssc	stc	sbc
uus	uds	ucs	uss	uts	ubs	sus	sds	scs	sss	sts	sbs
uut	udt	uct	ust	utt	ubt	sut	sdt	sct	sst	stt	sbt
uub	udb	ucb	usb	utb	ubb	sub	sdb	scb	ssb	stb	sbb
duu	ddu	dcu	dsu	dtu	dbu	tuu	tdu	tcu	tsu	ttu	tbu
dud	ddd	dcd	dud	dtu	dbd	tud	tdd	tcd	tsd	ttu	tbd
duc	ddc	dcc	dsc	dte	dbc	tuc	tde	tcc	tsc	tte	tbc
dus	dds	dcs	dss	dtu	dbu	tus	tds	tcs	tss	ttu	tbs
dut	ddt	dct	dst	dtu	dbt	tut	tdt	tct	tst	ttu	tbt
dub	ddb	dcb	dsb	dtb	dbb	tub	tdb	tcb	tsb	ttb	tbb
euu	edu	ecu	esu	etu	ebu	buu	bdu	bcu	bsu	btu	bbu
cud	cdd	ccd	csd	ctd	cbd	bud	bdd	bcd	bsd	btd	bbd
cuc	cdc	ccc	csc	cte	cbe	buc	bdc	bcc	bse	bte	bbc
cus	uds	ucs	uss	uts	ubs	bus	bds	bcs	bss	bts	bbs
cud	cdt	cct	cst	ctt	cbt	but	bdt	bct	bst	btt	bbt
cub	cdb	ccb	csb	ctb	cbb	bub	bdb	bcb	bsb	btb	bbb

216 possible Baryon topologies [excluding anti-matter] with each combination producing differing Baryonic nett energy levels

Quarks always form symmetrical [udu] or [dud] charged Baryonic topologies [same charges never combine except under high energy conditions] ie. Proton is a [udu] not [uud] as is commonly stated



Baryons

24
[30-06]



Charmed Sigma

Unstable High energy particle (extremely short lived)



12
[24-12]




Charmed Sigma




0
[18-18]



Charmed Sigma



12
[24-12]




Bottom Sigma



0
[18-18]




Bottom Sigma



12
[12-24]





Bottom Sigma



Unstable High energy particle (extremely short lived)


Baryons




Unstable High energy particle (extremely short lived)




Charmed Xi Prime




Charmed Xi Prime















Double Charmed Xi



Double Charmed Xi



Baryons

<p>0 [18-18]</p> 	<p>Bottom Xi (Cascade B)</p> 
<p>12 [12-24]</p> 	<p>Bottom Xi (Cascade B)</p> 
<p>0 [18-18]</p> 	<p>Double Bottom Xi</p> 
<p>12 [12-24]</p> 	<p>Double Bottom Xi</p> 
<p>12 [24-12]</p> 	<p>Charmed Bottom Xi</p> 
<p>0 [18-18]</p> 	<p>Charmed Bottom Xi</p> 

Tetryonics 10.13 - Charmed Baryons


Baryons

0

[18-18]



Charmed Omega




12

[12-24]



Bottom Omega




Unstable Highenergy particle (extremely short lived)

12

[24-12]



Double Charmed Omega




0

[18-18]



Charmed Bottom Omega




12

[12-24]



Double Bottom Omega




12

[24-12]



Double Charmed Bottom Omega




0

[18-18]



Charmed Double Bottom Omega



ZPF

Photons

Bosons


Tetryons

$n\pi$ charged mass-Energy-Matter geometries

Ω

ZPFs are the quantum of Bosons

Q



Υ

All energies seek equilibrium

Q

Energy momenta

EM mass-Energies

Charges

Matter

$$n\pi \left[\left[\frac{m\Omega}{\text{mass}} \frac{v}{\text{velocity}} \right]^2 \right]$$

$$\text{EVENT}\pi \left[\left[\frac{\epsilon_0 \mu_0}{\text{ElectroMagnetic}} \cdot \frac{m\Omega v^2}{\text{mass velocity}} \right] \right]$$


$$\text{ODD}\pi \left[\left[\frac{\epsilon_0 \mu_0}{\text{ElectroMagnetic}} \cdot \frac{m\Omega v^2}{\text{mass velocity}} \right] \right]$$

$$4n\pi \left[\left[\frac{\epsilon_0 \mu_0}{\text{ElectroMagnetic}} \cdot \frac{m\Omega v^2}{\text{mass velocity}} \right] \right]$$

Gluons


The gluon can be considered to be the fundamental exchange particle facilitating the strong interaction between protons and neutrons in a nucleus.

Gluons are the exchange particles for the 'colour' strong force between quarks, analogous to the exchange of photons in the electromagnetic force between two charged particles.





The interaction between neutral Tetryons (Gluons) and all charged Tetryons is the Strong Colour force

nett charge of Gluons is same as a Neutron





45,012
Electrons contain no Gluons



Ω


Gluons are considered to be bi-coloured, carrying a unit of colour and a unit of anti-colour

$\bar{\Omega}$



Q

Gluons are the neutral Tetryons inside all fermionic Matter topologies





They are the neutral charge [di-electric] Tetryons located between charged Tetryons in Matter topologies [attractive coulombic forces between their opposite charged fascias create the 'colour force']

Gluons in Fermions

In the Standard model Gluons have no mass and no electric charge. They carry a special 'colour charge' property that hold quarks together to form Baryons



Tetryonics reveals that gluons are neutral charge tetryons comprised of equal positive & negative mass-energies & are elementary Matter particles

'Gluons' are neutral tetryons



$$4\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

EM Field Planck quanta
Gluons ElectroMagnetic mass velocity



up Quarks have 1 Gluon
(Neutral Tetryon)


anti-down Quarks have 2 Gluons
(Neutral Tetryons)

GLUONS
 are the neutral tetryons in Matter topologies acting as a di-electric between charged tetryons

down Quarks have 2 Gluons
(Neutral Tetryons)


anti-up Quarks have 1 Gluon
(Neutral Tetryon)

Neutrinos are 3 Gluon sets



$$12\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$


EM Field Planck quanta
Glueballs ElectroMagnetic mass velocity




Baryonic Gluons

In 'the Standard Model', Gluons are vector gauge bosons that mediate strong interactions of quarks in quantum chromodynamics (QCD). Unlike the electrically neutral photon of quantum electrodynamics (QED), gluons themselves carry colour charge and therefore participate in the strong interaction in addition to mediating it, making QCD significantly harder to analyze than QED.

They are considered to be elementary particles which act as the exchange particles (or gauge bosons) for the colour force between quarks, analogous to the exchange of photons in the electromagnetic force between two charged particles



Deuterium nuclei (being the constituent quanta of all Elements) have 9 neutral Tetryons (Gluons) which in turn contribute to their gravitational mass along with the charged Tetryons



Tetryonics simplifies the current definition of Gluons and clearly identifies their geometric properties, along with their role in particle genesis

Neutrons have 5 Gluons

[Neutral Tetryons]

0

[18-18]



12


[24-12]

Protons have 4 Gluons


[Neutral Tetryons]

'Glueballs'

In particle physics, a glueball is a hypothetical composite particle. It solely consists of gluon particles, without valence quarks.




Neutral Tetryons are the result of equal number charged EM fields combining to form neutral charge Tetryonic topologies



It is conceivable that given the right conditions (ie a cloud of neutral Tetryons), that in the absence of charged Tetryons to interact with, a Glueball topology can be formed entirely from neutral Tetryons

Note: Despite their total neutral charge Glueball topologies are polarised

Neutrinos and some exotic neutral charge Baryons can be considered to be 'Gluon topologies'





Neutrons are not considered to be Glueballs as they contain charged Tetryons

OMEGA Particle Icosahedron Glueball




20 Neutral Tetryons




The 3 internalised planes formed by the external apex points correspond with the 3 spatial dimensions of Cartesian geometry

Non-gluonic Baryon formation


Non-neutral 'UP' quark has +4 charge as opposed to +8 charge of UP quark



8
[10-2]
+2/3




4
[4-8]
-1/3





Baryonic Particles can be formed with non-neutral dielectric Tetryons in lieu of the usual neutral Tetryons [Gluons] - altering the nett charges of the quarks formed

4
[8-4]
+1/3




4
[4-8]
-1/3







Un-stable form
[rapidly decays into constituent Quarks]



Resulting in another family of Baryons [CHARGEBALLS]

Stable Form
[possibly mistakenly viewed as Bottom Quarks]



Electro-Magnetic fields


Quantised Charge
is the nett geometry of mass-Energy-Matter formed by equilateral ElectroMagnetic fields and can be modelled with classical vector flux rotations



Positive Charge ZPF


Magnetic Monopoles do NOT exist

Negative Charge ZPF




Charge
All mass-Energy-Matter EM field interactions can be modelled using ZPF geometries

Current



The charge geometry of all EM fields determines their Electric Permittivity and Magnetic Permeability

The 'zero point' field has scalar EM energies



The EM field is a equilateral waveform with the Magnetic field always orthogonal to the Electric

The Magnetic fields propogate bi-directionally and the Electric field is responsible for producing linear momentum

The Electric field and Magnetic fields are equal to each other and directly proportional to the velocity of propagation

Photons
are the neutral quanta of EM waves they are comprised of two opposite charge EM field quanta




Zero Point fields are polarised and are the sources of



Electro-static and Magneto-static fields and particles

The Weak Force




6 Inductive 'Weak' edges
(3 internalised)
(3 externalised)

Quantum Energy is stored or released, and distributed throughout Tetryonic geometries in ODD number quanta [Bosons]




6 Inductive 'Weak' edges



Quantum Inductive loops
Quantum Tank circuit
Energy received is stored in quantum inductive loops (until its release via weak interaction - Boson exchange)

All ElectroMagnetic fields are ideal quantum inductive loops of EM energy momenta
Each has a Magnetic edge (base) which acts as a quantum inductor through which energy can be absorbed or released in EM quanta via electroMagnetic induction [exchange Bosons]

Leptons and Quarks have the same Tetryonic numbers but differing Matter topologies





Leptons and Quarks result from the Strong 'colour' force interactions in dodeca-deltahedrons

(In Quarks the attractive strong force produces Octahedral topologies)
 (In Leptons the repulsive strong force produces Dodecahedral topologies)



The Strong Force

The force resulting from the interaction of 2 Tetryons along their parallel planar (Electric charge) surfaces

'Attractive' Colour Force
Opposite charge Tetryon fascias

Same direction QAM flux


Colour Force fascias

Colour force Interaction angle


'Repulsive' Colour Force
Same charge Tetryon fascias

Opposing direction Electric flux rotation

Strong Force strength is directly proportional to the nett mass-energy quanta involved (Increases/decreases as total Tetryon mass-energy quanta increases/decreases)






The Strong Colour Force is the binding force that holds Nuclei together





Strong Force Interaction

On the smallest scale (less than about 0.8 fm), it is the Electric force (mediated by gluons in nuclei) that holds Tetryons together in order to form Quarks, Protons and Neutrons.



The residual Electric field force produced by the Strong Force is also facilitates the binding of Protons and Neutrons together to form the nucleus of an atom



Attractive Colour Charge Force holding Quarks and Nuclei together

Atomic Nucleus



36 charged mass-energy fascia geometries



12 [42-30]

creates an external Baryons topology with 20 charge fascia

Proton - Hydrogen ion

The Proton is a Baryon with a Positive Tetryonic charge geometry of +12 [24-12] [and a final Matter topology of 20π].

It attracts electrons [0-12] and Neutrons [13-18] through their opposite charged fascia in order to reach charge equilibrium

The Proton's opposite charge particle is the electron. If it binds with one or more Neutrons it will still attract electrons in order to achieve Tetryonic charge equilibrium.

As a Hydrogen nuclei it is highly reactive

Neutral Hydrogen

Hydrogen is the Second Neutral Tetryonic geometry (Baryon) formed in the creation of Matter

It is a [24-24] NEUTRAL [balanced] Tetryonic charge geometry

The Neutral Hydrogen Atom interacts with Deuterium and other Elements via its external nucleon fascia charges (residual EM Interactions)

Deuterium

Deuterium has the same nett charge as Neutral Hydrogen but has an increased Tetryonic mass- Matter topology

It has an additional 9 Tetryons in its nucleus (due to the Neutron) and consequently is larger than the Hydrogen nucleus

It has a [42-42] NEUTRAL Tetryonic charge geometry. Deuterium nuclei combine to form all other elements

Tritium

(Radioactive Deuterium)

Tritium has the same nett Tetryonic charge as Neutral Hydrogen and Deuterium (D) but due to its higher quantum energy levels it has additional mass-energy quanta in its charged geometry

It has the same 84 charged fascia in its nucleus as Deuterium but its increased mass-energy is equivalent to that of 1 Neutron.

This extra nuclear energy is the source of its radioactivity and ongoing confusion in chemistry where it is released as a variety of decay particles over time

Proton
Hydrogen ion

36

1 Proton	24-12
Total Charge	24-12 [+12]

H^+
F+ n=25




Neutral
Hydrogen

48

1 Proton	24-12
1 electron	0-12
Total Charge	24-24 [0]

H^0
F+ n=25
e- n=1




Deuterium

84

1 Proton	24-12
1 Neutron	18-18
1 electron	0-12
Total Charge	42-42 [0]

D
N+ n=25
F+ n=25
e- n=1




Tritium


84

1 Proton	24-12
1 Neutron	18-18
1 electron	0-12
Total Charge	42-42 [0]


D
N+ n=31
F+ n=31
e- n=7
Radioactive Isotope




Hydrogen - Helium genesis



Hydrogen Family



Proton 36π	electron 12π	Hydrogen 48π
----------------------	------------------------	------------------------



Hydrogen 48π	Neutron 36π	Deuterium 84π
------------------------	-----------------------	-------------------------




Deuterium 84π	Tritium 84π
-------------------------	-----------------------



Tetryonics 13.03 - Hydrogen family

Hydrogen-Helium3 genesis



Elemental Form





Positive Ions



Radioactive



Tritium 0
is a non-radioactive allotrope of Tritium,
it has the same Tetryonic charge and mass-Energy as Helium 3
(but is ionically 1 elemental charge less than Helium 3)




Quantum Batteries

Atomic nuclei can be easily scaled to non-quantum sizes to offer clean, safe and portable long term energy storage devices that can store energy indefinitely and release it on demand anywhere in the World




12 loop quantum inductive rotor

The quantum battery is unique in that in addition to storing energy indefinitely, when the nuclei combine with a lepton it has the ability to release specific energies [photons] by way of its synchronous quantum convertor topology



Energy stored in quantum batteries is measured as mass

(Atomic Nuclei)




anti-Parallel Configuration



The non-neutral charge of atomic nuclei attract free leptons into 'bound' states within the various n levels of atomic shells releasing energy as spectral photons

Series Configuration



Quantum synchronous Converters

With the addition of a quantum rotor (lepton) a quantum battery can be converted from a storage device into a energy distribution device.


And just like the quantum battery, the quantum convertor can be scaled to any size in order to provide tailor-made energy efficient delivery devices

The electron can be viewed as a rotating inductor consisting of 3 negative Tetryons

$$12\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]^{n8}$$

Leptons ElectroMagnetic mass velocity

The electron has a charged Matter topology that is electrically equivalent to a quantum 6 loop inductive rotor



$$72\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]^{n32}$$

Nuclei ElectroMagnetic mass velocity

$$84\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]^{n32}$$

Deuterons ElectroMagnetic mass velocity

Where varying levels and frequencies of Energy are transmitted long distances and need to be stored for later release on demand the 'ideal' mechanical device is the rotating (or synchronous) convertor

Changes in energy-momenta results in photon emission/absorpt on lines photon emission/absorption produces changes in leptonic energy-QAM and results in the quantum transition of electrons in atomic orbitals


Changes in Baryonic energy levels induces a directly proportional change in electron energy levels

3 forms of mass-energy momenta are possessed by quantum converters

Angular Momentum (motional energy)
linear momentum of Baryons & leptons

Photonic spectral energies (emission/absorption)
KEM fields geometries of photo-electrons


Nuclear mass-energies (stored energy quanta)
standing wave mass-energies of Matter topologies



A change in any 1 of the 3 types of energy in a atom results in a proportional change in the other 2


External EM fields and incident photons can all affect the quantum energy levels of the atomic nuclei

$$Mv^2 = KEM = hv^2$$





Hydrogen Ionisation Energies

The ionisation energy level for each quantum level is proportional to the square of the quantum number




In Hydrogen nuclei electrons remain in the same position close to the Proton due to Coulombic attraction.

The established model of electron orbitals having circular radii of increasing size in proportion to their energies is incorrect



Hydrogen Energy Levels

KEM field energy in each electron's energy level



Kinematics

An inertial frame of reference is one in which the motion of a particle not subject to forces and results in motion in a straight line at constant velocity

Sir Issac Newton



(25 December 1642 – 20 March 1727)

$$a = \frac{\Delta y}{\Delta x} = \frac{\Delta v}{\Delta t}$$

$$\mathbf{F} = ma \rightarrow \mathbf{a} = \mathbf{F}/m$$

Acceleration

[Inertial Frame PLUS odd quanta]



Deceleration

[Inertial Frame MINUS odd quanta]



Newton's First Law

$$\sum \mathbf{F} = 0 \Rightarrow \frac{d\mathbf{v}}{dt} = 0$$


Every body persists in its state of being at rest or of moving uniformly straight forward, except insofar as it is compelled to change its state by force impressed

PHILOSOPHIÆ
NATURALIS
PRINCIPIA
MATHEMATICA.

AUCTORE
ISAACO NEWTONO, EQ. AVX.

Ëditio tertia aucta & emendata.

LONDINI:
Apud GULIELMUM & JOHANNES ISSACUM, Regiæ Societatis typographos.
MDCCLXXVI.



Newton's Second Law

$$\sum \mathbf{F} = \frac{d\mathbf{p}}{dt} = m \frac{d\mathbf{v}_{cm}}{dt} = m\mathbf{a}_{cm}$$

Force creates a change in Momentum over Time

A body of [m]ass subject to a net [F]orce undergoes an [a]cceleration that has the same direction as the force and a magnitude that is directly proportional to the force and inversely proportional to the mass, i.e., $F = ma$.


Alternatively, the total force applied on a body is equal to the time derivative of linear momentum of the body.

Inertia and Force

momentum

The classical definition of Momentum relates the mass of a material body at given velocity (v) to its Momentum (p); it is a proportionality factor in the formula


$$p = mv$$



Linear momentum is the nett square root of mass-energy quanta

p

Any change to the energy-momenta content of a closed inductive loop requires a proportional change to the loop's energy density



[all inductive loops resist changes to their energy levels]

Inertia is the resistance of any physical object to a change in its state of motion or rest, or the tendency of an object to resist any change in its motion.

The principle of inertia is one of the fundamental principles of classical physics which is used to describe the motion of matter and how it is affected by applied forces


$$E = mv^2$$

Any change in the energy-momenta content of a body of mass-Matter results in a proportional change in its momentum-velocity

Force

This meaning of a body's inertia therefore is altered from the classical definition of "a tendency to maintain momentum" to a description of the measure of how difficult it is to change the momentum of a body

$$F = ma$$



Force is the sum of the linear mass-energy momenta quanta

F

Gottfried Wilhelm von Leibniz




(July 1, 1646 – November 14, 1716)

Scalar Energy

Leibniz's vis viva (Latin for living force) is mv^2 , twice the modern kinetic energy. He realized that the total energy would be conserved in certain mechanical systems, so he considered it an innate motive characteristic of Matter.

His thinking gave rise to another regrettable nationalistic dispute. His vis viva was seen as rivaling the conservation of linear momentum championed by Newton in England and by Descartes in France; hence academics in those countries tended to neglect Leibniz's idea.

In reality, both energy and momentum are conserved, so the two approaches are equally valid.



The nett direction of Force within energy geometry is UNIDIRECTIONAL ie the force exerted is the result of the nett linear momenta irrespective of charge

mass x velocity squared $kg \cdot \left[\frac{m}{s}\right]^2$



Energy $kg \cdot \frac{m^2}{s^2}$

$$J = \frac{kg \cdot m^2}{s^2} = N \cdot m = Pa \cdot m^3 = W \cdot s$$

A Joule is equal to the energy expended (or work done) in applying a force of one newton through a distance of one metre (1 newton metre or N·m)

momentum x velocity $\left[kg \cdot \frac{m}{s}\right] \cdot \frac{m}{s}$

Planck's quanta per second $\left[kg \cdot \frac{m^2}{s}\right] \cdot s^{-1}$



Linear Momentum

The Energy-momentum relationship is a fundamental physical property used to determine the mass of a body

Using the formula for mass-Energy equivalence as it relates to Photons moving at 'c'

$$E = h\nu = \frac{hc}{\lambda} = mc^2$$

Noting that the rest mass in the case of EM fields (Bosons and Photons) is to equal Zero

we can derive a relationship for Momentum - Energy - Wavelength showing that

$$p = E/c$$


Thus momentum in Photons is directly related to the EM energy content of the photon and the mass-energy content of any massive 3D body

$$p = \hbar k = \frac{h\nu}{c} = \frac{h}{\lambda}$$

and momentum in Particles is related to the total EM Energies of an object (its rest Matter + KE) and the wavelengths associated with those distinct energy levels

$$p = \frac{E}{c} = \frac{h\nu}{c} = \frac{h}{\lambda}$$

Linear momentum is the SQUARE ROOT of equilateral Planck mass-energy geometries and produces a unidirectional vector force



Longitudinal [linear] momentum

$$p = n\pi \left[\left[\begin{matrix} \text{Planck quanta} \\ m \Omega v \\ \text{mass} \quad \text{velocity} \end{matrix} \right] \right]_{\text{momentum}}$$

Momentum is a conserved quantity, meaning that the total momentum of any closed system (one not affected by external forces) cannot change.

Linear Momentum is the intrinsic square root vector component of Force

$$p = mv$$

In classical mechanics, momentum is the product of the mass and velocity of an object.

$$\sum \mathbf{F} = \frac{d\mathbf{p}}{dt} = m \frac{d\mathbf{v}}{dt} + \mathbf{v} \frac{dm}{dt}$$

In relativistic mechanics, this quantity is multiplied by the Lorentz factor.

$$p = n\pi \left[\frac{\left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \\ \text{mass} \quad \text{velocity} \end{matrix} \right]}{v} \right] = mv$$

EM field momentum is a function of its energy density, and is directly proportional to the group velocity

$$\left[\sqrt{\left[\begin{matrix} \text{Planck quanta} \\ m \Omega v \\ \text{mass} \quad \text{velocity} \end{matrix} \right]^2} \right]$$

Vector linear momentum

is the square root of the KEM field energies produced by Matter in motion

Classical formulation

$$\mathbf{p} = m\mathbf{v}$$

Linear Momentum is the vector force of mass times its velocity




$$\frac{E}{c^2} \cdot v$$

$$n\pi \left[\left[\begin{array}{c} \text{Energy} \\ m\Omega v \\ \text{momenta} \end{array} \right] \right]$$

Quantum formulation

$$\mathbf{p} = \frac{E}{v}$$

Linear Momentum is Energy divided by the square root of its quanta



$$\mathbf{p} = \sum_{i=1}^n m_i \mathbf{v}_i = m_1 \mathbf{v}_1 + m_2 \mathbf{v}_2 + m_3 \mathbf{v}_3 + \dots + m_n \mathbf{v}_n,$$

The linear momentum of a EM field or system of particles is the vector sum of the linear momenta in the KEM fields of individual particles in any spatial co-ordinate system



$$\frac{hv^2}{v}$$

Kinetic Energy

is the Electric field energy of Matter in motion

$$\frac{1}{2} \left[4n\pi \left[\epsilon_0 \mu_0 \cdot \left[m \Omega v^2 \right] \right] \right]$$

EM Field
Planck quanta
Matter
ElectroMagnetic mass velocity

Kinetic Energy is the diamond electric field extending from charged Matter topologies as a result of its motion, it follows Tetronic omega geometry and is proportional to an object's mass-Matter and its vector velocity

Charge
Q
[v-v]

Relativistic mass-Energies
M+KE

Kinetic Energy is 1/2 of the secondary KEM field energies created when Matter particles are in motion




$$2\pi \left[\frac{m \Omega v^2}{c^2} \right]$$

Planck quanta
mass velocity
c²
 EM Field

and is shown to be subject to Lorentz corrections

12n
rest Matter

[n-n]
Kinetic Energies



Kinetic Energies create
Magnetic Moments

$$KE = \frac{1}{2} M v^2$$

Kinetic energy is a scalar quantity; it does not have a direction.


The Kinetic Energy of an object is the energy which it possesses due to its motion.

It consists of Neutral Electric fields and an associated Magnetic moment

Kinetic Energies are subject to Lorentz corrections

$$E_k = mc^2 - m_0c^2$$

rest Matter is not



Kinetic Energy and Magnetic moments

$$Mv^2 = KEM = p^2$$

Unlike charged fields KEM fields have neutralised Electric Fields



KEM field geometry of positive charged Matter in motion




KEM field geometry of negative charged Matter in motion

$$KE = \frac{1}{2}Mv^2$$

Often noted as being two distinct EM energies [Kinetic Energy and Magnetic moments] are shown to be orthogonal aspects of the same KEM field of motion

As the velocity of a particle increases so does its Kinetic Energy and Magnetic moment creating Lorentz variable KEM fields



$$\mu_B = \frac{1}{2}Mv^2$$



$$\mu_B = \frac{1}{2}Mv^2$$

Types of Momentum



quantised angular momenta

[m²/s]

The quantised 'angular momentum' of each Planck mass-energy geometry, gives rise to the two quantum Charges



$$h$$

[kg-m²/s]

$$m\Omega$$



There are 3 forms of momenta in physics


equilateral charge geometry



Linear momentum

[kg-m/s]


The square root (v) of each Planck quanta's mass-energy geometry (v^2) is vector Linear Momentum



vector momentum

$$mv$$

[kg-m/s]




Angular momentum

[kg-m²/s]

The orbital angular momentum of electrons in atoms associated with a given quantum state

... [n-1] [n] [n+1] ...

vector rotation about a point




$$L^2 = L_x^2 + L_y^2 + L_z^2$$



Kinetic Energy vs Momentum

An important difference is that Kinetic energy is a scalar quantity - it has no direction in space
 momentum is a vector quantity - it has a direction in space, momenta combine like forces do.

$$p = Mv$$



$$KE = 1/2 [Mv^2]$$



$$p = \sqrt{E}$$

$$KEM = Mv^2$$


In Tetryonic geometry, the square root maps the linear momenta [mv] of a field to its ENERGY

$$p_{/2m}^2 = KE$$

In Tetryonic geometry, E-field geometry maps the kinetic energy [1/2 mv] of a field to its ENERGY

Linear momentum has a different geometric ENERGY relationship to that of *Electric fields*

mass-Energy-momentum




Lorentz velocity correction Factor


The Lorentz factor or Lorentz term is an expression which appears in several equations in special relativity. It arises from deriving the Lorentz transformations. The name originates from its earlier appearance in Lorentzian electrodynamics - named after the Dutch physicist Hendrik Lorentz.

Wavelength, momentum

mass-ENERGY




$$\beta = \left[\frac{v}{c} \right]$$



$$\left[\frac{v^2}{c^2} \right] = \beta^2$$

*Classically modelled as an infinite series approaching c
Tetryonics reveals it to be a physical property
of equilateral energy-momenta geometries*




**A vector measure of the
Energy content of
a waveform**

**A scalar measure of a
KEM waveform's
energies**

$$\gamma \equiv \frac{c}{\sqrt{c^2 - v^2}} = \frac{1}{\sqrt{1 - \beta^2}} = \frac{dt}{d\tau}$$

$$\gamma = 1/\sqrt{1 - v^2/c^2}$$

γ



Importantly - Newton's classical velocity addition remains correct, but the energy-momenta required to accelerate Matter to higher velocities increases exponentially in line with the equilateral [Tetryonic] geometry of mass-energies

Γ

Unified Energy momenta geometry

$$m = n\pi \left[\left[\epsilon_0 \mu_0 \right] \left[m \Omega v^2 \right] \right]$$

mass Electromagnetic mass Planck quanta velocity



$$p = n\pi \left[m \Omega v \right]$$


energy Planck quanta mass velocity



$$E = n\pi \left[\left[m \Omega v \right]^2 \right]$$



$$mv = p \qquad p^2 = E$$





$$\beta = \left[\frac{v}{c} \right]$$

Linear
Lorentz factor

$$\beta^2 = \left[\frac{v^2}{c^2} \right]$$

Scalar
Lorentz factor




$$E_\gamma = 2\pi \left[\left[\epsilon_0 \mu_0 \right] \left[m \Omega v^2 \right] \right]$$

Photons Electromagnetic mass Planck quanta velocity

$$m = 4n\pi \left[\left[\epsilon_0 \mu_0 \right] \left[m \Omega v^2 \right] \right]$$

mass Electromagnetic mass Planck quanta velocity

Squared Energy distributions



Bosons


Transverse EM mass-Energies

Odd number sequence

Number of quanta per geometry

$$n = (2n-1)$$

$h\nu$



EM mass-energy

Quantum numbers are *not* TRIANGULAR NUMBERS

$$\text{TRI } n = (n/2) \times (n+1)$$

Quantum numbers are *not* TETRAHEDRAL NUMBERS

$$\text{TETRA } n = (n/6) (n+1) (n+2)$$

Scalar energy levels have SQUARED quanta

Photons

Longitudinal EM mass-Energies

Even number sequence


Number of quanta per geometry

$$n = (2n)$$

Tetryonic Energy and Charge relationships



Quantised angular momenta is the geometric source of all mass-Energy-Matter relationships



The Energy density of a particle's charged fascia geometry determines its mass

m

Charged mass-energy geometry & Matter topology determines a particle's physical characteristics (Type, Family, generation, mass etc)


M

Charge

-12

electron

12
[0-12]



1.2 e20

mass-Matter

e-

8.851486 e-31 KG

Opposite nett elementary charge geometries

+12

Proton

12
[24-12]




2.25 e23

1875x
(differential)


differing Matter topologies

1.65965 e-27 KG

P+



Planck's energy momenta quantum



Planck mass **m** m - mass quantum

Planck Charge **Ω** v - quanta of Charge
quantum angular momentum spatial co-ordinate system
elementary charge / 12

Planck Energy **E** E - Energy quantum

v² Compton Frequency **mc² / h** 1×10^{19} quanta per second

λ de Broglie wavelength **h / mc** $2.99792458 \times 10^{-11}$ meters

p Momentum **mc** 2.21134×10^{-28} Kg

Planck's constant is the equilateral geometry of rest mass-energy momenta in standing wave Matter topologies

Currently the Atomic masses are calculated using 1/12 of a Molar Carbon12 atom's mass as the reference mass.

Tq [Carbon] - [252-252] **C** [270,072] / 12
 Tm [Carbon] - 270,072 **12** 22,506

If we calculated for the molar mass of C12 in 12 grams we would get
6.0121842700723 $7.378205107 \times 10^{-12}$ kg/mol
 this is in error by 1/2 of the mass of an electron



Tetryonic mass [Hydrogen] - 2.2512 e23 v

Using Tetryonic geometry we can solve for an exact Compton frequency of any mass-Matter topology and account exactly for all quantum energy contributions to the rest Matter of all electrons and Baryons in any element thus determining exact molar masses exclusive of blackbody radiation, kinetic energies and energies of measurement and avoiding 'weighted' masses

Defining [N] as the number of rest mass Hydrogen atoms in 1 gram exactly and makes Avagadro's number the inverse of this number

Molar mass = $\frac{H_1 \text{ Atomic mass}}{\text{Avagadro's No.}}$

mass H_{mol}/Av = $\frac{.001}{6.022141579 \times 10^{23}}$

rest mass Hydrogen = $1.660538841 \times 10^{-27}$ g/mol


H_{mole} / m[H] = $\frac{1.660738412 \times 10^{-27}}{22,512}$

n Planck mass = $7.376238634 \times 10^{-32}$ Kg


This is an exact Atomic rest mass as opposed to the current weighted molar Atomic mass estimates (which is incorrect by 1/2 the mass of an electron)

Planck's constant is the 'quantum of action' and also contributes to the mass-energies of radiant EM wave geometries


Nuclear Energy levels



Exponential energy levels




$$1e19v = \hbar = 5e18f$$



$$T\pi \left[\left[\frac{Energy}{mass} \Omega \right] \right]$$

Radioactive Decays follow exponential curves determined by the Tetryonic topology of the particle families





- Proton
- Neutron
- antiNeutron
- antiProton

- Electron
- Positron

- Neutrino

- Up
- Down
- Strange
- Charmed
- Top
- Bottom



- Positive
- Negative
- Neutral

- Bosons


$$ODD\pi \left[\left[\frac{EM\ Field}{ElectroMagnetic} \cdot \left[\frac{Planck\ quanta}{mass} \cdot \left[\frac{m\Omega v^2}{velocity} \right] \right] \right] \right]$$

Bosons

$$2\pi \left[\left[\frac{EM\ Field}{ElectroMagnetic} \cdot \left[\frac{Planck\ quanta}{mass} \cdot \left[\frac{m\Omega v^2}{velocity} \right] \right] \right] \right]$$

Photons

Linear energy levels



- 1 [1-0]
- 0 [1-1]
- 1 [0-1]

Emission and Absorption of Bosons and Photons within Atomic Nuclei

Increase and decrease in integer amounts according to the Tetryonic geometries and topologies of the particles involved

- 12 [2-4-12]
- 0 [8-18]
- 12 [12-24]
- 12 [12-0]
- 12 [0-12]
- 0 [6-6]
- 8 [10-2]
- 4 [4-8]
- 4 [4-0]
- 4 [2-2]
- 4 [0-4]


Energy momenta geometry

m

All mass-energies have equilateral geometries

E


mass-Energy
 $\text{kg} \frac{\text{m}^2}{\text{s}^2}$



Quantised angular momenta facilitates EM interactions

h

Planck's Constant
 $\text{kg} \frac{\text{m}^2}{\text{s}}$



scalar mass velocity squared


$$mv^2 = E = hv^2$$



quantised angular momenta per second

m
mass

E
Energy



Bosons are transverse EM masses

Photons are longitudinal EM masses

mass is a measure of Energy per spatial co-ordinate system

mass-Energy momenta

Electromagnetic energy is a scalar measurement of mass


**Bosons are
Transverse masses**



Planck mass

$$\text{ODD } \pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Bosons ElectroMagnetic mass velocity



**mass-ENERGY-Matter
equivalence**


**Photons are
Longitudinal masses**



Photons

$$\text{EVEN } \pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$


Photons ElectroMagnetic mass velocity



Tetryonic Matter of Electron - $8.851486361 \times 10^{-31}$ Kg
mass-Energy of Electron - $7.955319207 \times 10^{-14}$ J

$$12 \pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Fermions ElectroMagnetic mass velocity




Tetryonic Matter of Proton - $1.659653693 \times 10^{-27}$ Kg
mass-Energy of Proton - $1.491622351 \times 10^{-10}$ J

$$36 \pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Baryons ElectroMagnetic mass velocity

ELEMENTS



Tetryonic Matter of Deuterium - $3.320192534 \times 10^{-27}$ Kg
mass-Energy of Deuterium - $2.984040234 \times 10^{-10}$ J

$$84 \pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Deuterons ElectroMagnetic mass velocity

EM mass quanta in Matter

KG

kg




q

$$n\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

mass



f



$$ODD\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Bosons

ODDπ

EVENπ

$$EVEN\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

EM waves

Q



3 m

4 m

M

Matter

$$4n\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Matter



$$\beta = m \left[\frac{v}{c} \right]$$

[Linear momentum corrections are linear]

Lorentz velocity corrections

[result from the measurement of EM mass-energy quanta in a planar spatial co-ordinate system]

$$\beta^2 = m \left[\frac{v^2}{c^2} \right]$$

[EM mass-Energy corrections are Scalar]

$$m \quad \frac{E}{c^2} = \frac{\overset{\text{Energy}}{[m\Omega v^2]}}{\underset{\text{EM Field}}{c^2}} = \underset{\text{mass}}{n\pi} \left[\underset{\text{ElectroMagnetic}}{[\epsilon_0 \mu_0]} \cdot \overset{\text{planck quanta}}{[m\Omega v^2]} \right] \quad \left[\frac{1}{c^2} \right] E$$

$$p \quad mv = \left[\frac{E}{v^2} \right] \cdot v = \underset{\text{linear momentum}}{n\pi} \left[\underset{\text{ElectroMagnetic}}{[\epsilon_0 \mu_0]} \cdot \overset{\text{EM Field Planck quanta}}{[mv]} \right] \quad \frac{E}{v}$$




$$E \quad mv^2 = \frac{mv^2}{c^2} = \underset{\text{EM Field}}{4n\pi} \left[\underset{\text{ElectroMagnetic}}{[\epsilon_0 \mu_0]} \cdot \overset{\text{Energy}}{[mc^2]} \right] \quad m \left[\frac{c^2}{c^2} \right]$$

mass-energy in Matter propagates at c

$$KE \quad \frac{1}{2} Mv^2 = \frac{1}{2} \left[4 \left[\frac{\overset{\text{Energy}}{[m\Omega v^2]}}{\underset{\text{EM Field}}{c^2}} \right] \right] = \underset{\text{Photons}}{2\pi} \left[\underset{\text{ElectroMagnetic}}{[\epsilon_0 \mu_0]} \cdot \overset{\text{planck quanta}}{[m\Omega v^2]} \right] \quad 2m \left[\frac{v^2}{c^2} \right]$$




The geometry of Constants


Quantised angular momentum is the source of all physical Constants



A physical constant is a physical quantity that is generally believed to be both universal in nature and constant in time.

It can be contrasted with a mathematical constant, which is a fixed numerical value but does not directly involve any physical measurement





Max Planck




(April 23, 1858 – October 4, 1947)

"To interpret the vibrational energy of N oscillators not as a continuous, infinitely divisible quantity, but as a discrete quantity composed of an integral number of finite equal parts."

$$\epsilon = nh\nu$$

"Let us call each such part of the energy element h"



Planck's Constant

[quantised mass-energy angular momenta]

N_A

1 mole of Hydrogen atoms has a rest-mass of 1 gram

Solving for Planck's Constant using the inverse of Avogadro's number & Tetryonic geometry we obtain an exact corrected value of:

$$+ 6.629432672 \times 10^{-34} \text{ J.s}$$

$$- 4.137664516 \times 10^{-15} \text{ eV.s}$$

$$mv^2 = E = hv^2$$

A rest mass Hydrogen atom has a Compton frequency of 2.2512 e23 Planck quanta

0 [24-24]



2.2512 e23

Hydrogen

e-

P+



$$T\pi \left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \text{Matter} & \left[\epsilon_0 \mu_0 \cdot [m \Omega v^2] \right] \\ \text{ElectroMagnetic} & \text{mass} & \text{velocity} \end{matrix} \right]$$

Planck's constant is the relationship between EM mass-energy and quantised angular momenta that provides the basis for EM charge in Tetryonic geometries



(9 August 1776 – 9 July 1856)

Avogadro's Number

The number of rest mass Hydrogen atoms in 1 gram (and the rest molar mass of any element or compound) can be determined directly from tetryonic theory

[exclusive of any measurement, blackbody or kinetic energies]

Using a Compton frequency of 2.2512 e23 Planck quanta for a rest mass-Matter Hydrogen atom




atomic mass unit

$$1 \text{ u} = \frac{M_u}{N_A} = 1.660\,538\,782(83) \times 10^{-24} \text{ g}$$

$$\text{Avogadro } N = 6.022141579 \times 10^{23}$$

$$N^{-1} = \frac{2.2512 \text{ e}23 \text{ v}}{1.660538841 \times 10^{-24} \text{ g}} \text{ [Hydrogen mass]}$$



22,512
Hydrogen

n0 $1 \text{ mol} = 1 \text{ g}$
Hydrogen = $1.660538841 \times 10^{-27} \text{ kg}$

n1 $1 \text{ mol} = 1.000533 \text{ g}$

n0 $1 \text{ mol} = 11.996801 \text{ g}$
Carbon 12 = $1.99211552 \times 10^{-26} \text{ kg}$

n1 $1 \text{ mol} = 12 \text{ g}$

Hydrogen's rest Tetryonic mass is 22,512 n Planck quanta [Proton - 22,500n + electron 12n]

$$\left[22,512 \left[\left[\begin{matrix} \text{EM Field} \\ \epsilon_0 \mu_0 \end{matrix} \right] \cdot \left[\begin{matrix} \text{Planck quanta} \\ m \Omega v^2 \end{matrix} \right] \right] \right]^{-1}$$

ElectroMagnetic mass velocity

The inverse mass of Hydrogen is equal to Avogadro's number


Coulomb's Constant

$$k = \frac{1}{4\pi\epsilon_0}$$


The proportionality constant k_e , called the Coulomb constant (sometimes called the Coulomb force constant), is related to defined properties of EM Energy-momenta and is used to define Electric field forces

$$c = \frac{1}{\sqrt{\mu_0\epsilon_0}}$$


$$8.987 \text{ e9 } \frac{\text{Nm}^2}{\text{C}^2}$$



Similar repel



Opposites attract




Similar repel

Linear Coulombic force interactions are a result of charged E field linear momenta

$$\mathbf{E} = \frac{\mathbf{F}}{q_t}$$

The Electric field can be defined as the longitudinal Force exerted by charged masses



It is a measure of the interactive force produced by the Electric field energy-momenta of two superpositioned charge KEM fields

Longitudinal E field forces between Charged particles are mediated by Photons

$$\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \hat{\mathbf{r}}$$

The Electric field can also be derived from Coulomb's Law


C

The Speed of ElectroMagnetic energies

The classical behaviour of the electromagnetic field is described by Maxwell's equations, which predict that the speed c with which electromagnetic waves (such as light) propagate through the vacuum is related to the electric constant ϵ_0 and the magnetic constant μ_0 by the equation $c = 1/\sqrt{\epsilon_0\mu_0}$.

The speed of EM energies in a vacuum is defined as 299,792,458 meters per second (1,079,252,848.8 km/h).

Celeritas is a Latin word for "swiftness" or "speed". It is often given as the origin of the symbol c , the universal notation for the speed of light in a vacuum



The distance energy travels in 1 second from its source

EM waves and energy momenta can be measured as either Transverse or Longitudinal waveforms with respect to their velocity vector


All EM waves and energy are symmetrical waveforms whose quanta contain ElectroMagnetic mass-energy and momenta

The equilateral geometry of EM energy is the inverse of the spatial geometry of 1 second

The square root of equilateral energies [scalar speed] is a vector velocity

velocity is a vector direction of the energies of motion

speed is a scalar property of the energies of motion



The maximum velocity possible due to electrical accelerations or the impedance of any spatial region that energy propagates through it is often referenced in physics as 'the speed of Light' IT IS NOT AN ABSOLUTE LIMIT or BARRIER

All EM waves radiate outwards in accord with Biot-Savart's law

The velocity vectors of EM energies form the geometries of spatial co-ordinate systems

$$\frac{K_E}{K_M} = 9e16 \frac{Nm^2}{c^2} \frac{Ns^2}{c^2}$$

$$C^2 = 9e16 \frac{m^2}{s^2}$$

$$C = 299,792,458 \frac{m}{s}$$

The natural velocity of EM energies can be calculated from the field's EM permittivity & permeability and is affected by the medium it is propagating through

EM waveforms are bidirectional radiant emissions of 2D energy




with velocity [Lorentz] variable contents

mass quanta



EM mass-energy quanta

Matter is a KEM standing wave propagating at c



Permittivity x Energy density

$$\frac{E}{c^2} = m$$



$$\text{mass} = 7.376238634 \times 10^{-51} \text{ Kg}$$

Planck-Einstein

Quantum masses

$$h\nu^2 = E = m\nu^2$$

$\text{kg} \frac{\text{m}^2}{\text{s}} \frac{1}{\text{s}} \quad \text{kg} \frac{\text{m}^2}{\text{s}^2} \quad \text{kg} \frac{\text{m}^2}{\text{s}^2}$

The quantum of mass-energy can be derived with several methodologies using Tetryonic Geometry

Planck mass-energy geometries

quantised energy momenta

$$h$$

$$\text{kg} \frac{\text{m}^2}{\text{s}}$$

6.629432672 e-34 J

Planck quanta

$$[m\Omega v^2]$$

mass velocity

7.376238634 e-51 kg

$$\text{kg} \frac{\text{m}^2}{\text{s}^2}$$

$$m$$

scalar mass-energy

Charged mass-Matter topologies


The mass of Matter

$$\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

ElectroMagnetic mass velocity

Matter Quanta

Tetryonic molar mass [Hydrogen] - 1 g



$$\text{Molar mass} = \frac{H_i \text{ Atomic mass}}{\text{Avagadro's No.}}$$

$$\text{mass H}_{\text{mol}} / \text{Av} = \frac{.001}{6.022141579 \times 10^{23}}$$

$$= 1.660538841 \times 10^{-27} \text{ g/mol}$$

$$H_{\text{mole}} / \Delta m [H] = \frac{1.660538841 \times 10^{-27}}{22,512}$$

$$\text{mass} = 7.376238634 \times 10^{-32} \text{ KG}$$

Avogadro - Mandeleev


2D mass-Energy equivalence

The Speed of Light is the Natural velocity of Energy propagation

Energy

$$mv^2 = hv^2$$

EM mass



Zero Point Fields
 EM Field Planck quanta
 $n\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$
 Quanta ElectroMagnetic mass velocity




$m=1\pi$

mass-energy

[1, 2, 3, 4, 5, 6, 7, 8, 9, 10.....]


radiant 4π mass-energy geometries



in standing wave topologies
are 3D MATTER quanta

[2D EM masses are planar energy momenta]

Photons
 EM Field Planck quanta
 $2n\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$
 Photons ElectroMagnetic mass velocity



$m=2\pi$

EM energy carriers

[2, 8, 18, 32, 50, 72, 98, 128.....]

The geometry of mass and Matter

[1, 3, 5, 7, 9, 11, 13, 15, 17, 19.....]

[3D Matter has mass-energy momenta and volume]

[4, 8, 12, 24, 36, 48, 72, 84, 168.....]


Charge Carriers



$m=3\pi$


Bosons
 EM Field Planck quanta
 $ODDn\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$
 Bosons ElectroMagnetic mass velocity

Bosons



3D Matter topologies are standing waves of
2D mass-energies propagating at c

$m=4\pi$




mass-Matter
 EM Field Planck quanta
 $4n\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$
 Matter ElectroMagnetic mass velocity

mass-Matter

mass geometries & rest Matter topology

2D
 m
 mass
 kg



M
 Matter
 KG
 3D

Quanta number

$1e19 v$

Compton frequency

Matter is a three dimensional charged mass-energy topology

$$M = 4n\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$


EM Field
Planck quanta
Matter
ElectroMagnetic
mass
velocity

Quantum number

n

Nuclear Level

Energy density is mass - the term 'massless particle' is a misnomer




Kinetic Energies

$$RE = \text{Matter} + KE$$

Relativistic mass is the total EM energy content of a massive body (or system) in motion

The relativistic rest mass-energies of Matter are velocity invariant as 3D Matter is a EM standing wave with 2D mass-energy fascia whose velocity of propagation is the speed of Light



Matter

Charge topologies and rest mass-energies

mass-Matter topologies

mass-energy geometries

12
[0-12]

Electron 1.2×10^{-20}

n=0
Q
0.00000053 g
molar mass

Tm Electron = 12n

electron Charge	$-1.602216081 \times 10^{-19} \text{ C}$
electron rest mass	$8.851486361 \times 10^{-31} \text{ kg}$
electron rest energy	$7.955319207 \times 10^{-14} \text{ J}$ 496,519.7 eV (ELECTRON MASS)

12
[24-12]

Proton 2.25×10^{-23}

n=25²
Q
0.000999 g
molar mass

Tm Proton = 22,500n

Proton Charge	$+1.602216081 \times 10^{-19} \text{ C}$
Proton rest mass	$1.659653693 \times 10^{-27} \text{ kg}$
Proton rest energy	$1.491622351 \times 10^{-10} \text{ J}$ 930,974,522.8 eV (PROTON MASS)

Protons and Neutrons have identical masses to each other and equivalent topologies but differing charges

0
[18-18]

Neutron 2.25×10^{-23}

n=25²
Q
0.000999 g
molar mass

Tm Neutron = 22,500n

Neutron Charge	0
Neutron rest mass	$1.659653693 \times 10^{-27} \text{ kg}$
Neutron rest energy	$1.491622351 \times 10^{-10} \text{ J}$ 930,974,522.8 eV (NEUTRON MASS)

Hydrogen

When Protons and electrons interact to form neutral Hydrogen the electron's KEM quantum field energies increase in direct proportion to the Proton's energy levels

0
[24-24]

Hydrogen 2.2512×10^{-23}

n=0
Q
1.000533 g
molar mass

Tm Hydrogen = 22,512n

Hydrogen = Proton + electron

H Charge	0
H rest mass	$1.660538841 \times 10^{-27} \text{ kg}$
H rest energy	$1.492417883 \times 10^{-10} \text{ J}$ 931,967,562.3 eV (HYDROGEN MASS)

0
[42-42]

Deuterium 4.5012×10^{-23}

quantum synchronous converters

n=0
Q
1.999466 g
molar mass

Tm Deuterium = 22,512n

Hydrogen = Proton + electron

H Charge	0
H rest mass	$3.320192534 \times 10^{-27} \text{ kg}$
H rest energy	$2.984040234 \times 10^{-10} \text{ J}$ 1,862,445,565 eV (DEUTERIUM MASS)

In order to make an exact 1kg reference mass-Matter topology



$$\frac{\text{mass}}{c^2} \left[\frac{\text{Planck quanta}}{m \Omega v^2} \right]$$

mass-energy
Energy per second

$$mc^2 = E = hv^2$$

$$m = E/c^2$$

mass is the scalar integral surface area of Matter topologies

which is equivalent to exactly 1.000 molar Hydrogen masses

1,000 N_A

3.2512e23

whose Tetryonic Matter topology can be modelled using charged Planck mass-energy geometries

grams per kg

1,355704512 e50

6.629432672 e-34 J

Energy


8.021747579 e23

Avogadro's number

3.2512e23

1kg

revealing that there are exactly 1,355704512 e50 Planck quanta in standing wave Matter



$$\frac{\text{Matter}}{c^4} \left[\frac{\text{Planck quanta}}{m \Omega v^2} \right]$$

Matter-energy
Energy per second squared

$$hv^2 = E = Mc^4$$

$$M = E/c^4$$



Matter topologies are standing-waves of mass-energy

you need to create a standing-EM wave with a specific compton frequency


EM mass and Matter defined

There remains a lot of confusion over the exact definition of EM mass and Matter resulting in the frequent interchanging of one term for the other in physical processes

This must be clarified and the two terms must be properly defined in a manner that explains their derivation and physical properties in detail.



EM mass is a measure of equilateral scalar energy per unit of Time



Matter is the 3D topology of standing wave 2D mass-energies

EM mass
3D Matter is comprised of 2D mass-energies
2D mass-energy cannot contain 3D Matter
Matter

$\beta = \left[\frac{v}{c} \right]$
mass is the Lorentzian velocity corrected energy content of Tetryonic geometries

All forms of EM mass-Energy are subject to Lorentz factor relationships

$\beta^2 = \left[\frac{v}{c} \right]^2$
Matter is Lorentz invariant as its mass-energies propagate in a 4π standing wave geometry



$$M = \frac{4\pi E}{c^4}$$

Bosons and Photons are not 'massless' they are 'Matterless' [2D waveforms]

Bosons

$$m_p = \frac{h\nu}{c^2}$$


Planck masses



Photons

$$m_\gamma = \frac{hf}{c^2}$$

Photon mass



$$\text{ODD } \pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$


Bosons ElectroMagnetic mass velocity

$$\text{EVEN } \pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

EM waves ElectroMagnetic mass velocity



Charged mass-Matter geometries



Any measurement of a system's mass is subject to velocity corrections

mass = EM energy density

$$n\pi \left[\frac{\text{Planck quanta mass} \cdot \Omega \cdot \text{velocity}^2}{\text{Spatial geometry } c^2} \right]$$

n [7.376238634 e-51] kg

EM mass is a measure of 2D planar energies comprising the fascia of charged topologies

Matter Topology is determined by the geometry of charged Planck $T[q]$ quanta


ENERGY $n\pi \left[\frac{\text{Planck quanta mass} \cdot \Omega \cdot \text{velocity}^2}{c^2} \right]$ M

The mass - ENERGY - Matter contents of any physical system are all related through the spatial co-ordinate system used (which in turn is determined by the speed of light)



2D space c^2 3D space c^4

mass $\frac{n\pi}{c^2} \left[\frac{\text{Planck quanta mass} \cdot \Omega \cdot \text{velocity}^2}{c^2} \right]$ E Matter $\frac{T\pi}{c^4} \left[\frac{\text{Planck quanta mass} \cdot \Omega \cdot \text{velocity}^2}{c^4} \right]$

Matter and Charge are velocity invariant properties




As the energy content [levels] of particles and fields increase their intrinsic Planck quanta and EM mass increases but their charge geometries remain the same



Charged mass-Energies

All 3D Matter particles are comprised of charged fascia whose energy content determines their 2D mass



3D Matter geometries

Tetryons
4 charge fascia

Quarks
12 charge fascia

Neutrinos
12 charge fascia

Leptons
12 charge fascia

Baryons
36 charge fascia

Deuterium ion
72 charge fascia

$$\frac{\text{Matter}}{c^4} \left[\frac{\text{Planck quanta}}{[m \Omega v^2]} \right]$$

mass velocity

4π $2.651773069 \times 10^{-33} \text{ J}$
 $2.950495454 \times 10^{-50} \text{ kg}$
 4 quanta


12π $4.97207450 \times 10^{-11} \text{ J}$
 $5.532178976 \times 10^{-28} \text{ kg}$
 7.5×10^{22} quanta

12π $7.955319207 \times 10^{-33} \text{ J}$
 $8.851486361 \times 10^{-50} \text{ kg}$
 12 quanta

12π $7.955319207 \times 10^{-14} \text{ J}$
 $8.851486361 \times 10^{-31} \text{ kg}$
 1.2×10^{20} quanta

36π $1.491622351 \times 10^{-10} \text{ J}$
 $1.659653693 \times 10^{-27} \text{ kg}$
 2.250×10^{23} quanta

72π $2.984040234 \times 10^{-10} \text{ J}$
 $3.320192534 \times 10^{-27} \text{ kg}$
 4.5012×10^{23} quanta




In physics there exists
2 forms of energy momenta
[linear momenta &
quantised angular]

MOMENTA



The nett scalar
energy momenta
required to effect
changes in velocity

ENERGY



Net Charge is reflective of the total energy momenta
of any EM field or Matter subjected to acceleration
[inertial mass]


Inertial resistance to Force

Any change in motion results in changes to
the Charge geometries creating in turn proportional
changes to the nett KEM mass-energy momenta components

$$F = ma$$

The inductive resistance of quanta in charged EM fields
to any changes to their nett mass-energy geometry
is the source of what we term inertial mass

Inertia is the resistance of any physical object
to a change in its state of motion.



Force does work when it results in movement

$$W = F \cdot d = ma \cdot d = E$$

$$\text{kg} \frac{\text{m}}{\text{s}^2} \text{m}$$

classical mechanics

$$\Delta m v = \Delta E = \Delta m \Omega v$$

linear momenta

$$\text{kg} \frac{\text{m}^2}{\text{s}^2}$$

quantum mechanics

angular momenta

The Planck constant is the quantum of Action

$$E = m \Omega v^2 = h v^2$$

$$\text{kg} \frac{\text{m}^2}{\text{s}} \frac{1}{\text{s}}$$

Changing an object's **velocity** results in a corresponding **energy momenta** change which relate to each other through its **inertial mass**


EM field densities

EM mass is a measure of the energy content of any spatial co-ordinate system

$$\beta^2 = \left[\frac{v}{c} \right]^2$$

and is subject to Lorentz velocity corrections in 2D EM fields

6.629432672 e-34 J
 1 planck quantum has a EM mass-Energy of
7.376238634 e-51 kg
 and
 Quantised Angular Momentum which creates Charge
1.33518 e-20 C



mass velocity

mass-Energy equivalence



7.376238634 e-51 kg


$$mv^2 = E = hv^2$$

8.987551737 e16 [m/s]² 6.629432672 e-34 j quanta/sec

Matter is only mass-energy in Tetrahedral topologies [T4π+] [else it is nπ EM mass-energies that propagate away at c]

planck quanta







If reduced to a flat Euclidean space geometry Matter topologies become radiant mass-energies

m mass	E ENERGY	M Matter
$\left[\frac{n\pi}{c^2} \rho \right]$	$\left[m \Omega v^2 \right]$ <small>Planck quanta</small> <small>mass velocity</small>	$\left[\frac{4n\pi}{c^4} \rho \right]$

The inertial properties of electromagnetic mass - ENERGY & Matter [can all be differentiated as energy densities per unit of time] in any spatial co-ordinate system



m
[odd π]hv



1π Charge
EM Field Planck quanta
ElectroMagnetic mass velocity
 $[[\epsilon_0\mu_0] \cdot [m\Omega v^2]]$

Bosons

2D

Photons

2π Photons
EM Field Planck quanta
ElectroMagnetic mass velocity
 $[[\epsilon_0\mu_0] \cdot [m\Omega v^2]]$

2D planar Euclidean electromagnetic mass-energies propagate through the vacuum energy aether without interaction

$n\pi$ mass
EM Field Planck quanta
ElectroMagnetic mass velocity
 $[[\epsilon_0\mu_0] \cdot [m\Omega v^2]]$

m
[even π]hv




EM Field geometries & Matter topologies

**2D
Fields**

**3D
Matter**

Tetryonic charged geometries provides a clear mechanism for providing all particles with their distinct properties of 2D inertial EM mass-energy and 3D Matter.

M
[4n π]hv



12π Dodecyon
EM Field Planck quanta
ElectroMagnetic mass velocity
 $[[\epsilon_0\mu_0] \cdot [m\Omega v^2]]$

Leptons

3D


Baryons

36π Baryons
EM Field Planck quanta
ElectroMagnetic mass velocity
 $[[\epsilon_0\mu_0] \cdot [m\Omega v^2]]$

$T\pi$ Matter
EM Field Planck quanta
ElectroMagnetic mass velocity
 $[[\epsilon_0\mu_0] \cdot [m\Omega v^2]]$


3D standing wave topologies of electromagnetic mass-energies interact with the vacuum energy aether at various angles through their charged [inductive] fascia

M
[4n π]hv




Velocity invariant rest Matter

Electrostatic charges have no Magnetic Moments



All Matter are 3D standing wave topologies

$$M = \frac{E}{c^4}$$



$$E = Mc^4$$

rest mass-Matter is composed of $4\pi\pi$ standing wave topologies and is INVARIANT to velocity changes


The Relativistic EM mass-energies of a system in motion is the sum of its invariant rest Matter and Kinetic Energies

$$RE = \underbrace{12\pi}_{\text{Fermions}} \left[\frac{m\Omega v^2}{c^4} \right] + \underbrace{2\pi}_{\text{Photons}} \left[\frac{m\Omega v^2}{c^2} \right]$$

EM fields resulting from motion are subject to Lorentz correction


Kinetic mass-energies are divergent from invariant rest Matter topologies as a result of a particle's motion

At zero velocity the relativistic mass is equal to the invariant mass.



The energy which an object has due to its motion will not add mass into the invariant rest Matter of the particle in motion (it increases the total Planck quanta [EM mass-energies] of its extended KEM field)

Wavelengths are proportional to EM energy content



Photons are bidirectional Kinetic EMFields


$$\frac{E}{c^2} = m$$

Matter in motion has a resultant velocity related Energy field that possesses the physical properties of Kinetic energy and Magnetic moments

$$KEM = Mv^2$$

KEM field Energy is directly related to the Velocity changes of massive particles

All 2D EM fields and 3D Matter particles have ElectroMagnetic fields, inertial mass-energies & momenta resulting from their constituent equilateral Planck quanta which possess the additional physical properties of Compton Frequency and De Broglie Wavelength



EM mass is a measure of the inertial energy density of any 2D [EM field] or 3D [Matter] energy geometry

$$m_{\Omega} v^2 = E = m v^2$$

EM mass-energy momenta

The EM mass of an object is a fundamental property of the object; a numerical measure of its inertia; a fundamental measure of the energy density of an object.

mass geometries and Matter Topologies

All 3D Matter topologies contain 2D EM mass-energy geometries
not all 2D EM mass-energy geometries form 3D Matter topologies

EM mass can be clearly defined as a measure of the energy density of any charged geometry

$$T\pi \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m_{\Omega} v^2 \right] \right]$$

Labels: EM Field, Planck quanta, Matter, ElectroMagnetic, mass, velocity

3D Matter is any mass-energy geometry that creates a closed volume Topology (4π Tetryonic geometry)



$$\frac{E}{c^2} = m$$

EM waves [FIELDS] are distinguishable from Material Particles [MATTER] through their non-Tetrahedral topologies



mass-energy is a conserved property
Matter is not conservative




$$M = \frac{E}{c^4}$$

The term 'massless' is a misnomer and should be discontinued in its use as all EM fields and Particles have EM mass [Energy quanta per second] geometries [other alternatives could be 2D, EM field, or Matterless]




EM mass

$$\frac{E}{c^2} = m$$



$$M = \frac{E}{c^4}$$



Collapsed Matter

All attempts to clearly differentiate between planar EM mass-energy & 3D Matter in physics have met with limited success until the introduction of charged geometries.


mv mass-velocity squared is Energy



Planck quanta is Energy content **mΩ**

mass-ENERGY-Matter equivalence

Newton and Leibniz measured Energy as a velocity related function of mass



EM mass quantum

$$\frac{n\pi}{c^2} \left[\left[\underset{\text{mass}}{m\Omega} \underset{\text{velocity}}{v^2} \right] \right]$$

Energy momenta


$$mv^2 = E = hv^2$$

EM masses are subject to velocity related Lorentz corrections.



Matter is a standing-wave of electromagnetic energies and is Lorentz invariant

Planck and Einstein viewed mass-energies as quantised particles with momenta



Matter quantum

$$\frac{T\pi}{c^4} \left[\left[\underset{\text{mass}}{m\Omega} \underset{\text{velocity}}{v^2} \right] \right]$$

$$n\pi \left[\left[\underset{\text{mass}}{m\Omega} \underset{\text{velocity}}{v^2} \right] \right]$$

m

mass geometries

$$\frac{n\pi}{c^2} \left[\left[\begin{matrix} \text{mass} & \text{Planck quanta} \\ m & \Omega v^2 \end{matrix} \right] \right]$$

All forms of mass-ENERGY-Matter
are defined by their charged geometries and
the spatial co-ordinate systems used to measure their physics

M

Matter topologies

$$\frac{T\pi}{c^4} \left[\left[\begin{matrix} \text{Matter} & \text{Planck quanta} \\ m & \Omega v^2 \end{matrix} \right] \right]$$

Unified Field Equation

for Tetryonic EM mass-Energy-Matter

$$T\pi \left[\left[\begin{matrix} \text{Tetryonic} & \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & \cdot & m \Omega v^2 \end{matrix} \right] \right]$$

geometry ElectroMagnetic mass velocity

$$\frac{n\pi}{\text{mass}} \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$$

bosons & photons

nπ

EM waveforms

$$\frac{ODD\pi}{\text{Bosons}} \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$$

$$\frac{2\pi}{\text{Photons}} \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$$

$$\frac{EVEN\pi}{\text{EM waves}} \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$$



spatial co-ordinate systems
are defined by the velocity of energy

Fermions

Tπ

Particles

$$\frac{4\pi}{\text{Tetryons}} \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$$

$$\frac{12\pi}{\text{Dodecyons}} \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$$

$$\frac{24\pi}{\text{Mesons}} \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$$


$$\frac{36\pi}{\text{Baryons}} \left[\left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & m \Omega v^2 \end{matrix} \right] \right]$$

EM mass

Matter

m

M



mass
 $\frac{n\pi}{c^2} \left[\left[\frac{\text{Planck quanta}}{m} \Omega v^2 \right] \right]$
mass velocity

Planck quanta
 $n\pi \left[\frac{m \Omega v}{\text{mass velocity}} \right]^2$
 energy

Matter
 $\frac{T\pi}{c^4} \left[\left[\frac{\text{Planck quanta}}{m} \Omega v^2 \right] \right]$
mass velocity

Tetryonic unified field Geometry

p

mass velocity



7,376238634 e-51 kg

$$mv^2 = E = hv^2$$

6.629432672e-34 js

8.987551787 e16 [m/s]² 6.629432672 e-34 j quanta/sec

mass-Energy momenta equivalence



planck quanta

h

The equilateral geometry of quantised angular momentum is the key to understanding Energy

$\text{kg} \frac{m}{s}$

mv



$m\Omega$

$\text{kg} \frac{m^2}{s}$

linear momentum



charge

quantised angular momentum


Classical Mechanics

Quantum Mechanics


EM mass Relationships



EM mass is revealed to be the scalar property of 2D Energy waveforms that is at the core of many important physical processes and measurements



E/second



E/second²



2D mass geometries should never be confused with 3D Matter topologies; nor should the terms be used in exchange for each other

Rest Matter

$$\frac{\pi}{\text{mass-Matter}} \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

3D rest mass-Matter topology = closed volume of 2D mass-energies

3D Electrostatic particle
No Magnetic Moment



All Matter is a Tetronic standing-wave charged geometry occupying a volume in 3D spherical space

$$\left[\frac{\pi}{\text{Matter}} \left[\left[\frac{\text{Planck quanta}}{c^4} \right] \left[m \Omega v^2 \right] \right] \right]$$


↓

$$\left[\frac{12\pi}{\text{Dodecyon}} \left[\left[\frac{\text{Planck quanta}}{c^4} \right] \left[m \Omega v^2 \right] \right] \right]$$

rest mass-Matter is velocity invariant (not subject to Lorentz corrections)

Relativistic mass-Matter

The property of Matter cannot be measured using a planar [c squared] spatial co-ordinate system



A KEM field has Electric and Magnetic quanta whose total energy quanta is directly related to the square of the particle's velocity


$$4\pi \left[\frac{\text{mass velocity}}{c^2} \left[m \Omega v^2 \right] \right]$$

All KEM fields are subject to Lorentz corrections

Kinetic Energies

$$\frac{4n\pi}{\text{KEM field}} \left[\left[\epsilon_0 \mu_0 \right] \cdot \left[m \Omega v^2 \right] \right]$$

Matter-Energy



Relativistic mass Energy = rest Matter + Kinetic EM Energies

$$\left[\frac{12\pi}{\text{Lepton}} \left[\left[\frac{\text{Planck quanta}}{c^4} \right] \left[m \Omega v^2 \right] \right] \right] + \left[\frac{4\pi}{\text{KEM wave}} \left[\left[\frac{\text{Planck quanta}}{c^2} \right] \left[m \Omega v^2 \right] \right] \right]$$

Kinetic Energies

KEM mass-energies are velocity dependent (subject to Lorentz corrections)

m

Tetryonic mass & Matter

Historically interchanged due to the lack of proper definitions the physical properties of EM mass & Matter can now be firmly defined with respect to their energy equivalence and spatial geometries


M

mass is a measure of the 2D planar energy content of any physical system

Matter is a measure of the 3D volumetric energy content of any physical system


$$\frac{n\pi}{c^2} \left[\left[m \Omega v^2 \right] \right]$$

mass Planck quanta
mass velocity



$$\frac{T\pi}{c^4} \left[\left[m \Omega v^2 \right] \right]$$

Matter Planck quanta
mass velocity



EM mass should replace the generic term mass with reference to ElectroMagnetic energy densities



MATTER is a geometric 4π standing wave topology of EM mass-energy geometries

E/second

E/second²


2D
radiant equilateral geometry
of EM mass-ENERGY momenta
EM mass

3D
standing-wave topology
of EM mass-ENERGY momenta
rest Matter

The electromagnetic energies of rest Matter is never 'at rest' as the electromagnetic field energies creating mass-Matter topologies always propagate at c

Photoelectron KEM fields


The linear momenta produced as a result of energy momenta in KEM fields is converted into angular momentum when leptons are bound to atomic nuclei



The EM mass-energy content of Baryons directly influence the KEM field energy levels of bound Leptons




EM mass-energy-momenta



and Spin are always conserved

The motion of leptons within atomic nuclei produces Magnetic moments at various orientations to the nuclear magneton





If an electron is 'ejected' from the Nuclei it will obey conservation of EM mass-energy momentum

ie Its ejected energy-momentum equals the absorbed photon's energy-momenta [minus the work energy required to free it]




Nuclear magneton





Leptronic Oscillations



Antimatter

$$12\pi \left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \left[\epsilon_0 \mu_0 \right] & \left[m \Omega v^2 \right] \end{matrix} \right]$$

Leptons ElectroMagnetic mass velocity

All Leptronic generations, oscillations and types can be accounted for through Tetryonic geometries & energy levels

Matter

Eigenstate bound energy states




In order to ionise any photo-electron from its bound position within a Hydrogen atom the KEM field energy of the electron must be increased from its Eigenstate value to more than 13.525ev

Principal Quantum levels

$$Mv^2 = KEM = hv^2$$


Bound electron Eigenstate values



The mass-energy content of Baryons comprising atomic nuclei directly determines the Kinetic energy levels of bound Leptons [forming quantum synchronous convertors] (Incident Bosons/Photons can increase these levels if they contain the exact quanta required to increase the level to the next square number)


Bohr radius

As the Electron's angular velocity increases the measurable Bohr magnetic moment strength increases and its associated KEM field Planck wavelengths decrease (due to the increased Planck quanta).



The highest quantum number level produces the strongest Magnetic Moment

All massive particles absorb and release energy in discrete quantum steps according to their respective Tetryonic geometries and changes in velocities







Kinetic Energies







Nuclear Magnetons

are weaker than Bohr magnetons due to differing mass-charge ratios and non-parallel Quark magnetons



Baryons

are tri-Dodecyon geometries



$$36\pi \left[\begin{matrix} \text{EM Field} & \text{Planck quanta} \\ \epsilon_0 \mu_0 & [m \Omega v^2] \\ \text{Baryons} & \text{ElectroMagnetic mass velocity} \end{matrix} \right]$$

Tetryonic Charge

P^+

12
[24-12]
Q
36

Nuclear magneton

N^0

0
[18-18]
Q
36

Nuclear magneton

N^0

0
[18-18]
Q
36

Nuclear magneton

P^-

12
[12-24]
Q
36

Nuclear magneton

Matter geometry

Baryonic Magnetic Moments

0
[18-18]



N^0



P^+


12
[24-12]



A Baryon's Magnetic Moment
is a result of the combined non-parallel
Magnetic moments of its Quarks




0
[18-18]



N^0

P^+




12
[24-12]

As a result of the 3 non-aligned Quark magnetic moments
resulting from their Tetryonic geometry,
the Nuclear Magneton is considerably weaker than the Bohr Magneton

Nuclear Quantum Energies

[Principal Quantum Numbers]

All electron energy levels are reflective of the KEM field of a electron in a specific quantum level
(The rest Matter of each electron is invariant)




Higher Quantum Level electrons already have high Kinetic energies and thus require lower frequency photons (low additional KE) in order to be ejected from their bound nuclear positions

-13.6 eV Hydrogen ionisation Energy

An Electron's energy can only increase in steps that reflect the Tetryonic Matter geometry of Leptons and their square mass-energies

Energy required to ionise electron with an existing KEM field



Principal Quantum numbers reflect an electron's energy level


Energy may be absorbed or released from any lepton in Quantum steps reflecting the energy difference between the electron 'orbitals'

$$\frac{1}{\lambda} = \frac{R_H}{hc} \left[\frac{1}{n_i^2} - \frac{1}{n_f^2} \right]$$

Transitions between electron energy levels is the basis for emission and absorption spectra

$$KEM = \frac{E_i}{n^2} = \frac{-13.525 \text{ eV}}{n^2} \quad \text{Eigenstate energy levels}$$

Any electron that has in excess of 13.525 eV of Kinetic Energy has sufficient KE to escape the Nucleus



Leptons are historically classified as Spin 1/2 particles
(by the spin-statistics theorem and the Pauli exclusion principle)
as determined by their magnetic moments

Quantum Spin Numbers

(rotations about an axis)

Spin 1/2

720°

Rotating a spin-1/2 particle by 360 degrees does not bring it back to the same quantum state it needs a 720 degree rotation



Planck bar relates to the electric field content of KEM fields resulting from Matter in motion (as reference to the Nuclear magneton or external Magnetic field)

Spin 0

any°

A spin-zero particle can only have a single quantum state, even after torque is applied.



Spherical 'point particles' of charge do not exist



Spin 1

360°

Rotating a spin-1 particle 360 degrees can bring it back to the same quantum state



SPIN must not be confused with Chirality (reflections)



Spin 2

180°

Rotating a spin-2 particle 180 degrees can bring it back to the same quantum state



electro-static field



magneto-static field



electro-static field

Photons are their own anti-particle

Spin 3

120°


Rotating a spin-3 particle 120 degrees can bring it back to the same quantum state



On a geometric basis all Leptons are in fact spin 3 particles

Magnetic moments are determined by the KEM fields created by vector linear momentum

Nuclear Magnetic moments are complicated by the tri-quark magnetons within all Matter topologies



The KEM field of any charged particle in motion is reflective of the particle's nett charge topology



720°

Rotating a spin-1/2 particle by 360 degrees does not bring it back to the same quantum state it needs a 720 degree rotation

Spin 1/2

μ_B

Parallel

μ_N

anti-Parallel

μ_B

electron spins

μ_B

anti-Parallel

μ_N


Parallel

μ_B

120°

Rotating a spin-3 particle 120 degrees can bring it back to the same quantum state

Spin 3




All atomic particles have a particular "spin" analogous to the Earth's rotation on its axis.

$$\mu_B = \left[\frac{Q}{2mv} \right] / e$$

An isolated electron has an angular momentum and a magnetic moment resulting from its spin. While an electron's spin is sometimes visualized as a literal rotation about an axis, it is in fact a fundamentally different, quantum-mechanical phenomenon with no true analogue in classical physics.

The quantum mechanical reality underlying spin is complex and still poorly understood. Consequently, there is no reason to expect the above classical relation to hold.

$$\mu_B = g \left[\frac{e^-}{2mv} \right] S$$



Tetryonic KEM field geometries reveals the source and orientation of all atomic magnetic moments


Generating Magnetons

A static Electron has a negative Tetryonic charge [0-12] topology with neutralised magnetic dipoles



Leptons are 12 loop inductive charge rotors


Tetryonic geometry fully explains Leptronic 'spins'



Einstein's Special Relativity model of distorted moving charges producing magnetic moments is incorrect

BOHR Magnetron produced by Lorentzian distortion of charges due to relativistic velocities

$$\mu_B = \frac{e\hbar}{2m_e}$$




The term "electron spin" can now be taken literally (when modelled with Tetryonic geometries) as an accurate description of the origin of Magnetic moments for all Leptonic [BOHR] magnetons.

The previously held model of the electron as a spinning sphere of charge must be abandoned in favour of the true Tetryonic charge geometries of EM mass-Energy-Matter

A moving Electron has a KEM field with an Electric field and a Magnetic Moment



The gyromagnetic ratio of a particle or system is the ratio of its magnetic dipole moment to its angular momentum



The KEM field energy of an Electron in motion is subject to relativistic corrections due to energy changes resulting from its acceleration

$$\lambda = L' \sqrt{1 - \frac{v^2}{c^2}}$$

WAVE-length contraction of mass-energy quanta of KEM field

REPEL

Ampere Force

$$\mathbf{B} = \frac{\mu_0 I}{4\pi} \int \frac{d\ell \times \hat{r}}{r^2}$$

$$\mathbf{F} = 2k_A \frac{I_1 I_2}{r}$$


Lorentz Force

$$\mathbf{F} = q[\mathbf{E} + (\mathbf{v} \times \mathbf{B})]$$

ATTRACT

Stern Gerlach Experiment

In 1922 O. Stern and W. Gerlach measured the intrinsic spin angular momentum of silver atoms and found it to take only two discrete values, $+\hbar/2$ and $-\hbar/2$ commonly called "spin up" and "spin down"




The Stern-Gerlach experiment to determine electron SPIN

The results were interpreted to show that particles possess an intrinsic angular momentum that is most closely analogous to the angular momentum of a classically spinning charged sphere, that takes on only certain quantized values of angular momentum

Tetryonics shows the results are a product of the KEM field produced by charge particles in motion, with all charged particles being able to produce 2 distinct magneton orientations as a result of the real motion of their intrinsic quantum inductive loop (Matter) geometries through external E fields & as referenced to external M moments.

This along with the Lorentz force produced by an external Magnetic field acting on the KEM magnetons produces the two results obtained

Any increase in electron energies results in an increase in quantised angular momenta and its vector linear momentum




Negative charged in motion produce negative charge KEM field geometries

Any change in the KEM energy level of an electron results in Photon emission or absorption


Electron Spins based on the measured Bohr Magnetons of moving electrons are reflected with KEM field geometries

Electrons bound in atomic nuclei increase or decrease their energy levels dependent on the energy level of the nuclei in which they are bound

Bound and unbound electrons can rotate in one of two directions influenced only by external magnetic and electric fields or incident photons



The two separated beams of electrons produced are defined as having differing SPINS [UP or DOWN]




$$z = \frac{1}{2} a t^2 = \frac{1}{2} \frac{F}{m} \left[\frac{L}{v} \right]^2 = \pm \frac{\mu_B L^2}{4 KE} \frac{\partial B_z}{\partial z}$$

The orientation of the BOHR magneton wrt the NUCLEAR magneton determines the 'direction' of electron SPIN

Bohr Magnetons

A moving electron is a 12 loop rotating inductor

12
[0-12]



A moving Lepton creates a secondary stronger intrinsic magnetic dipole moment within its KEM field which interacts with external magnetic fields

12
[0-12]



Leptons are not point particles

Electro-static particles have neutralised Magnetic moments

Velocity creates Kinetic Energy and Magnetic moments

Left handed and right handed fermions are mirror images of each other

Energy created by moving a Lepton through an external EM field is stored as Planck quanta within the Lepton's extended KEM field

Reversing the vector direction of the particle's linear momentum creates a reversed dipole Magnetic moment

All Leptonic macro-KEM fields and interactions with external fields can be modelled using Tetryonic geometries



Leptonic 'spin' is always determined by the Leptonic Magnetic moment as referenced against the Nuclear Magnetic Moment




Magnetic moment 'spin orientations' are reversed for opposite charge particles

Electron Spin orientation

The Bohr magneton dipole produced by Kinetic Energies is located axially about the centre of rotation




All Leptons have 12 intrinsic neutralised dipole moments and a polarised KEM field Magnetic moment created by the energies of its motion



Lepton Magnetic moment

[Bohr Magnetron]

$$\mu_B = \frac{e\hbar}{2m_e}$$



Nuclear magnetons are weaker than Bohr magnetons

All Leptonic spin directions are referenced to external Magnetic fields [either Nuclear Magnetons or H fields]

Nuclear Spins

are determined by the orientation of Bohr Magnetons with respect to the Nuclear Magnetons [or an external Magnetic field]

electron Matter geometries are negative charged fascia which create neutral intrinsic magnetic dipole configurations





MAGNETIC MOMENT

In a moving electron Kinetic energy produces an axial Magnetic moment

In a static electron all intrinsic dipoles are neutralised through their orientations




Quantum Inductive Loops in motion produce Kinetic Energies in turn creating Magnetic Moments




Spin UP is a higher energy state resulting in a (Parallel) Bohr Magnetons

Nuclear magneton or external Magnetic field



Spin DOWN is a lower energy state resulting in a (antiParallel) Bohr Magnetons

Lorentz Force
When moving in an external magnetic field
The axial dipole moment of an electron will experience torque force proportional to its velocity
**Opposites attract
Similar repel**





Nuclear magnetic Moment

[Nuclear magneton]


NUCLEAR magnetons are much weaker than BOHR magnetons due to the higher charge to mass ratio of electrons

$$\frac{M_p}{M_e} = \frac{1}{1875}$$

The nuclear spins for individual protons and neutrons parallels the treatment of electron spin, with spin 1/2 and an associated magnetic moment.




For the combination of electron, neutrons and protons in periodic elements, the situation is even more complicated.



Spin orbital coupling mechanics

The induced magnetic moment of electrons in atomic nuclei combine vectorally with the magnetic moment of the nuclei


anti-parallel moments
Spin DOWN




μ_B
Bohr magnetons



parallel moments
Spin UP




The energy level differences created
are manifested in Hyperfine-line splitting
Zeeman effects etc.



Spin DOWN
anti-parallel moments

Nuclear magnetons
 μ_Z



Spin UP
parallel moments

The alignment of electron spins in nuclei results in diamagnetic, para-magnetic Matter

Gyromagnetic Ratio

The electron is a 12 charge quantum rotor with a uniform charge to mass density ratio, the ratio of its magnetic moment to its orbital angular momentum, also known as gyromagnetic ratio



This implies that a more massive assembly of charges spinning with the same angular momentum will have a proportionately weaker magnetic moment, compared to its lighter counterpart.



The Bohr Magnetron is determined by the charged KEM field geometry of Leptons

1875
charged mass-Matter differential

The combined Kinetic energy of Motion [KEM field] and Electron Spin coupling with Nuclear Magnetons will effect any measured Gyromagnetic ratios



Classical electron model
[rotating sphere]

9.1 e-31 kg

4π

An electron is NOT a spherical particle

12π

Tetryonic electron
[rotating tri-tetryon topology]

8.85 e-31 kg

In physics, the gyromagnetic ratio (also sometimes known as the magnetogyric ratio in other disciplines) of a particle or system is the ratio of its magnetic dipole moment to its angular momentum, and it is often denoted by the symbol γ , gamma.

An isolated electron has an angular momentum and a magnetic moment resulting from its spin.

Its SI units are radian per second per tesla (s⁻¹T⁻¹) or, equivalently, coulomb per kilogram (C·kg⁻¹).

Tetryonic quantum mass to Charge ratio

$$1.810109642 \times 10^{11}$$


The 2006 CODATA
- e / m_e = -1.758 820 150(44) x 10^[11]

Nuclear Magnetons

e ⁻	Tetryonic elementary charge 1.602216081 e-19 coulombs	12q
e _m	Tetryonic electron mass 8.851486361 e-31	1.2 e20

[q/m]


$$181,010,964,200 \text{ C/kg}$$



12q	Tetryonic elementary charge 1.602216081 e-19 coulombs	e ⁺
2.25 e23	Tetryonic Proton mass 1.659653693 e-27	P _m

[q/m]

$$96,539,180.9 \text{ C/kg}$$



Tetryonics 19.11 - DIA & PARA-magnetic fields

12
[12-0]



Positron

1.2e20

Leptons

181,010,964,200 C/kg

12
[0-12]



Electron

1.2e20

12
[24-12]



Proton

2.25e23

Baryons

96,539,180.9 C/kg

0
[18-18]



Neutron

2.25e23

96,487,720.78 C/kg

Elements

48,256,721.99 C/kg

0
[24-24]



Hydrogen


2.2512e23

Deuterium



4.5012e23

All mass-Matter have distinct charge geometries and once in motion are subject to EM forces as a result of their geometries & KEM field



Lorentz force Law

$$\vec{F} = q\vec{E} + q\vec{v} \times \vec{B}$$

LORENTZ force Electric force charge velocity Magnetic force

all Matter in motion is subject to NEWTON'S SECOND LAW OF MOTION

$$F = ma$$

Combining Lorentz's force law & Newton's Third Law we obtain a EM mass-charge quotient for any particle of mass-Matter in motion

$$\left(\frac{M}{Q}\right)a = E + v \times B$$


mass-charge ratios

1.335180067 e-20 C

$$1.810109642 \text{ e11} \left[\frac{q}{m}\right] n \text{ 1e19 V}$$

7.376238634 e-51 kg

are the basis for mass spectroscopy



Q Any two particles with the same EM mass-to-charge ratio **M** follow the same path in a vacuum when subjected to the same external electric field

Dependent on their generations


Charge/mass		
$\left(\frac{q_t}{m_t}\right)$	181,010,964,200 C/kg	$\left(\frac{q_e}{m_e}\right)$
4:4		12:12

Tetryons and Leptons can have identical mass-charge ratios


Tetryonic theory shows mass-charge ratios are a measure of mass-energy geometries in Matter topologies

Collider particle track physics

A collider is a type of a particle accelerator involving directed beams of particles.



Colliders may either be ring accelerators or linear accelerators, and may collide a single beam of elementary particles against a stationary target or two beams head-on.



2D bosons & photons are radiant mass-energy geometries

3D Matter Fermions are standing wave mass-energy topologies

Analysis of the by-products of these collisions without a clear definition of, and distinction between, EM mass-energy geometries & Matter topologies through the charged geometrics of Tetryons ceases a misleading and erroneous picture of the particles created in high energy collisions within particle physics accelerator experiments

E

Q

t 4
[0-4]

4 t
[4-0]

Tetryons have the same 1/3 elementary charge as some Quarks


d 4
[4-8]

4 d
[8-4]

m



181,010,964,200 C/Kg



Tetryons can have the same mass-charge ratios as Leptons




181,010,964,200 C/Kg



M

Radiant EM mass-energy geometries



Charge

$$\frac{1\pi}{c^2} \left[\frac{\text{Planck quanta}}{m \Omega v^2} \right]$$

Photons

$$\frac{2\pi}{c^2} \left[\frac{\text{Planck quanta}}{m \Omega v^2} \right]$$




2D Planar divergent EM mass-energies form radiant Electromagnetic waves


ElectroMagnetic mass is a property of Matter
[being the energy content of its charged geometry per unit time]

mass

$$\frac{n\pi}{c^2} \left[\frac{\text{Planck quanta}}{m \Omega v^2} \right]$$



2D space-time measurements of equilateral energies form the geometric basis for electromagnetic mass




Bosons

$$\frac{\text{ODD}\pi}{c^2} \left[\frac{\text{Planck quanta}}{m \Omega v^2} \right]$$

EM waves

$$\frac{\text{EVEN}\pi}{c^2} \left[\frac{\text{Planck quanta}}{m \Omega v^2} \right]$$



Q
charge




Quantised angular momenta
are energy geometries

Planck's constant of
mass-energy momenta
is the source of all physical
constants and force relationships

ENERGY

2D Equilateral scalar energy-momenta




ENERGY

$$n\pi \left[\left[\begin{matrix} \text{Planck quanta} \\ m\Omega v^2 \\ \text{mass} \quad \text{velocity} \end{matrix} \right] \right]$$

Equilateral energy-momenta form
the foundation of all Forces,
EM masses & Matter


All energy seeks equilibrium

E
energy




Squared numbers in physics
create equilateral geometries

Separated fields of charge
create electromotive forces
accelerating Material bodies
within them



Positive charge energy momenta fields




Negative charge energy momenta fields

3D Matter topology is NOT a property of 2D ElectroMagnetic mass-energies

it is a measure of the closed 3D standing-wave spatial topology of all fermionic particles created by their charged equilateral mass-energy momenta geometries

Matter displaces vacuum energies to create convergent gravity fields around it




Tetryons are the quantum of Matter

$$\frac{4\pi}{c^4} \left[\frac{\text{Planck quanta}}{\text{mass}} \left[m \Omega v^2 \right] \right]$$

$$\frac{12\pi}{c^4} \left[\frac{\text{Planck quanta}}{\text{mass}} \left[m \Omega v^2 \right] \right]$$

$$\frac{24\pi}{c^4} \left[\frac{\text{Planck quanta}}{\text{mass}} \left[m \Omega v^2 \right] \right]$$




3D Matter particles are Z component second squared mass-energy topologies & created closed volumes

$$\frac{\text{Matter}}{c^4} \left[\frac{\text{Planck quanta}}{\text{mass}} \left[m \Omega v^2 \right] \right]$$

The mass-energies of Matter are Lorentz invariant to velocity changes

All Matter is comprised of [and radiates divergent] kEM mass-energies



Deuterium is the quantum of all Elements


$$\frac{\text{Baryons}}{c^4} \left[\frac{\text{Planck quanta}}{\text{mass}} \left[m \Omega v^2 \right] \right]$$

$$\frac{\text{Nuclei}}{c^4} \left[\frac{\text{Planck quanta}}{\text{mass}} \left[m \Omega v^2 \right] \right]$$

$$\frac{\text{Deuterons}}{c^4} \left[\frac{\text{Planck quanta}}{\text{mass}} \left[m \Omega v^2 \right] \right]$$

Tetryonic Mechanics

The charged geometry of mass-ENERGY-Matter



ISBN 978-0-9872884-1-7
[Second Edition © 2012]

Abraham